



# Aurora A Kinase (Human) Assay/Inhibitor Screening Kit

## Product Information

---

### Cat.No.

Kit-0121

---

### Product Overview

Aurora A Kinase (Human) Assay/Inhibitor Screening Kit is a single-site, non-quantitative immunoassay for Aurora-A activity. Plates are pre-coated with a substrate corresponding to recombinant Lats2, which contains serine83 residues that can be phosphorylated by Aurora-A. The detector antibody specifically detects only the phosphorylated form of serine83 residue on Lats2.

---

### Description

Aurora-A is a key member of a closely related subgroup of serine/threonine kinases that belongs to the *Drosophila aurora* and *Saccharomyces cerevisiae* Ipl1 kinase family, and both are essential for chromosome segregation and centrosome functions. Aurora-A kinase has been shown to contribute to oncogenic transformation and is frequently overexpressed and amplified in many human tumor types. It has been reported that amplification of Aurora-A in approximately 12% of primary breast tumors, as well as in breast, ovarian, colon, prostate, neuroblastoma, and cervical cancer cell lines. Additionally, high expression of Aurora-A mRNA was detected in tumor cell lines without evidence of gene amplification. Ectopic expression of Aurora-A in mouse NIH 3T3 cells led to the appearance of abnormal centrosome number (amplification) and transformation in vitro. Finally, overexpression of Aurora-A in near-diploid human breast epithelial cells revealed similar centrosome abnormality, as well as induction of aneuploidy. These findings suggested that Aurora-A is a critical kinase-encoding gene, whose overexpression leads to centrosome amplification, chromosomal instability, and transformation in mammalian cells.

---

### Applications

1) Screening inhibitors or activators of Aurora-A. 2) Detecting the effects of pharmacological agents on Aurora-A activity.

---

### Target Species

Human

---



## Aurora A Kinase (Human) Assay/Inhibitor Screening Kit

### Usage

---

For research use only (RUO)

---

### Storage

---

Upon receipt store the ATP at -20°C Upon receipt store all other components at 4°C; Do not expose reagents to excessive light

---

### Kit Components

---

Microplate: One microplate supplied ready to use, with 96 wells (12 strips of 8-wells) in a foil, zip-lock bag with a desiccant pack. Wells are coated with recombinant Lats2-N-ter as an Aurora-A substrate. 10X Wash Buffer: One 100 mL bottle of 10X buffer containing 2% Tween -20. Kinase Buffer: One bottle containing 20 mL of 1X buffer; used for Kinase Reaction Buffer and sample dilution. 20X ATP: Lyophilized ATP Na<sub>2</sub> salt. Reconstitute contents of vial with 2 mL of H<sub>2</sub>O. Mix gently until dissolved. Final concentration of ATP should be 1 mM ATP. The ATP solution can be stored in small aliquots (e.g. 100 µL) at -20°C. The 1 mM ATP stock solution must be diluted to 50 µM in Kinase Reaction Buffer at the time of the assay. Anti-Phospho-Lats2-S83 Monoclonal Antibody (ST-3B11): One vial containing 12 mL of anti-phospho- Lats2-S83 monoclonal antibody (ST-3B11). Ready to use. HRP conjugated Anti-mouse IgG: One vial containing 12 mL of HRP (horseradish peroxidase) conjugated anti-mouse IgG. Ready to use. Substrate Reagent: 20 mL of the chromogenic substrate, tetra-methylbenzidine (TMB). Ready to use. Stop Solution: One bottle supplied ready to use, containing 20 mL of 1 N H<sub>2</sub>SO<sub>4</sub>.

---