

## TR-FRET FXR Coactivator Assay Kit, goat

### Product Information

---

**Cat**

Kit-0977

---

**Common Name**

TR-FRET

---

**Cat.No.**

Kit-0977

---

**Description**

The TR-FRET FXR (farnesoid X receptor) Coactivator Assay Kit provides a sensitive and robust method for high-throughput screening of potential FXR ligands as agonists of coactivator recruitment. The kit uses a terbium (Tb)-labeled anti-GST antibody, a fluorescein-labeled coactivator peptide, and an FXR ligand-binding domain (FXR-LBD) that is tagged with glutathione-S-transferase (GST), in a homogeneous, mix-and-read assay format.

To assay:

When running the TR-FRET FXR coactivator assay in agonist mode, FXR-LBD is added to ligand test compounds followed by addition of a mixture of the fluorescein-coactivator peptide and Tb-labeled anti-GST antibody. After an incubation period at room temperature, the 520 nm/495 nm TR-FRET ratio is calculated and used to determine the EC<sub>50</sub> from a dose response curve of the compound. Based on the biology of the FXR-coactivator peptide interaction, the ligand EC<sub>50</sub> is a composite value representing the amount of ligand required to bind to receptor, effect a conformational change, and recruit coactivator peptide.

---

**Applications**

Co-Factor Interaction Assay

---

**Storage**

The TR-FRET FXR Coactivator Assay Kit contains FXR-LBD (GST) protein, fluorescently labeled SRC2-2 coactivator peptide, Tb-labeled anti-GST antibody, and buffers. Store components as indicated in the assay protocol (-80°C, -20°C, or +4°C).



## TR-FRET FXR Coactivator Assay Kit, goat

### Shipping

---

Dry Ice

---

### Size

---

800 x 20  $\mu$ L assays

---

### Materials Required but Not Supplied

---

Microplate Reader

---

**Detection method** Fluorescent

---