



## TR-FRET FXR Coactivator Assay Kit, rabbit

### Product Information

#### Cat

Kit-0956

#### Common Name

TR-FRET

#### Cat.No.

Kit-0956

#### Description

The TR-FRET FXR (farnesoid X receptor) Coactivator Assay Kit provides a sensitive and robust method for high-throughput screening of potential FXR ligands as agonists of coactivator recruitment. The kit uses a terbium (Tb)-labeled anti-GST antibody, a fluorescein-labeled coactivator peptide, and an FXR ligand-binding domain (FXR-LBD) that is tagged with glutathione-S-transferase (GST), in a homogeneous, mix-and-read assay format.

To assay:

When running the TR-FRET FXR coactivator assay in agonist mode, FXR-LBD is added to ligand test compounds followed by addition of a mixture of the fluorescein-coactivator peptide and Tb-labeled anti-GST antibody. After an incubation period at room temperature, the 520 nm/495 nm TR-FRET ratio is calculated and used to determine the EC50 from a dose response curve of the compound. Based on the biology of the FXR-coactivator peptide interaction, the ligand EC50 is a composite value representing the amount of ligand required to bind to receptor, effect a conformational change, and recruit coactivator peptide.

#### Applications

Nuclear Receptor Assay

#### Storage

Tb-anti-GST Antibody (Rabbit): Store at -20 °C

FXR-LBD, GST: Store at -80 °C

Fluorescein Coactivator Peptide: Store at -20 °C



CREATIVE **BIOMART**<sup>®</sup>  
Assay Kit

## TR-FRET FXR Coactivator Assay Kit, rabbit

TR-FRET Coregulator Buffer G: Store at 4 °C

1M DTT: Store at -20 °C or -80 °C

---

### Size

800 x 20 µL assays

---

### Materials Required but Not Supplied

Microplate Reader

---

**Detection method** Fluorescent

---

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)

Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA