



IKK-alpha/beta Kinase (Human) Assay/Inhibitor Screening Assay Kit

Product Information

Cat.No.

Kit-0475

Product Overview

IKK-alpha/beta Kinase (Human) Assay/Inhibitor Screening Assay Kit is a single-site, non-quantitative immunoassay for activities of IKK-alpha and IKK-beta. Plates are pre-coated with a substrate corresponding to recombinant I κ B-alpha, which contains two serine residues that are phosphorylated by IKK-alpha and IKK-beta (I κ B kinases). The detector antibody specifically detects only the phosphorylated form of I κ B-alpha.

Description

Although a number of recent studies suggest that I κ B degradation and nuclear translocation of NF- κ B may not be the sole regulatory events in the transcription of NF- κ B-dependent genes, it has been proposed that there is the central dogma of NF- κ B activation, which suggests that NF- κ B is sequestered in the cytoplasm in resting cells by the inhibitory I κ B proteins. In response to a variety of agonists, I κ B is rapidly phosphorylated, ubiquitinated, and degraded, thus releasing NF- κ B for translocation into the nucleus to initiate gene transcription. I κ B kinase (IKK) is the convergence point in most signaling pathways activated by many stimuli leading to the inducible phosphorylation and degradation of I κ B. IKK is a multisubunit complex that contains two catalytic subunits, IKK α and IKK β ; and the regulatory subunit IKK γ . Gene knock out studies have clearly demonstrated that IKK β ; and IKK γ subunits of the IKK complex are required for NF- κ B activation by all known pro-inflammatory stimuli including lipopolysaccharide (LPS), TNF, and IL-1. Thus a selective inhibitor of IKK β ; would not only be of great interest as a potential anti-inflammatory agent but also as a valuable tool to understand the mechanisms regulating NF- κ B activation by these inflammatory agonists.

Applications

1) Screening inhibitors or activators of IKK α and IKK β ; 2) Detecting the effects of pharmacological agents on IKK α and IKK β ; activity.



IKK-alpha/beta Kinase (Human) Assay/Inhibitor Screening Assay Kit

Target Species

Human

Usage

For research use only (RUO)

Storage

• Upon receipt store the ATP at -20°C • Upon receipt store all other components at 4°C; Do not expose reagents to excessive light

Kit Components

Microplate: One microplate supplied ready to use, with 96 wells (12 strips of 8-wells) in a foil, zip-lock bag with a desiccant pack. Wells are coated with recombinant IκBa as substrate of IKK. 10X Wash Buffer: One 100 mL bottle of 10X buffer containing 2% Tween-20. Kinase Buffer: One bottle containing 20 mL of 1X buffer; used for Kinase Reaction Buffer and sample dilution. 20X ATP: Lyophilized ATP Na₂ salt. Reconstitute contents of vial with 1.6 mL of H₂O. Mix gently until dissolved. Final concentration of ATP should be 1.25 mM ATP. The ATP solution can be stored in small aliquots (e.g. 100 μL) at -20°C. The 1 mM ATP stock solution must be diluted to 62.5 μM in Kinase Reaction Buffer at the time of the assay. Anti-Phospho-IκBa S32 Monoclonal Antibody (AS-2E8): One vial containing 12 mL of anti-phospho-IκBa S32 monoclonal antibody (AS-2E8). Ready to use. HRP conjugated Anti-mouse IgG: One vial containing 12 mL of HRP (horseradish peroxidase) conjugated anti-mouse IgG. Ready to use. Substrate Reagent: 12 mL of the chromogenic substrate, tetra-methylbenzidine (TMB). Ready to use. Stop Solution: One bottle supplied ready to use, containing 12 mL of 0.5 N H₂SO₄.