



## TR-FRET TR beta Coactivator Assay Kit, goat

### Product Information

---

**Cat**

Kit-0896

---

**Common Name**

TR-FRET

---

**Cat.No.**

Kit-0896

---

**Description**

The TR-FRET TR (thyroid receptor) beta Coactivator Assay Kit provides a sensitive and robust method for high-throughput screening of potential TR beta ligands as agonists of ligand-dependent coactivator recruitment. The kit uses a terbium (Tb)-labeled anti-GST antibody, a fluorescein-labeled coactivator peptide, and a human recombinant TR beta ligand-binding domain (TR beta-LBD) that is tagged with glutathione-S-transferase (GST) in a homogeneous, mix-and-read assay format.

To Assay:

When performing the TR-FRET TR beta coactivator assay, TR beta-LBD is added to ligand test compounds followed by addition of a mixture of the fluorescein-coactivator peptide and Tb anti-GST antibody. After an incubation period at room temperature, the 520 nm/495 nm TR-FRET ratio is calculated and can be used to determine the EC<sub>50</sub> from a dose response curve of the compound. Based on the biology of the TR beta-coactivator peptide interaction, the ligand EC<sub>50</sub> is a composite value representing the amount of ligand required to bind to receptor, induce a conformational change, and recruit coactivator peptide.

---

**Applications**

Co-Factor Interaction Assay

---

**Storage**

Store in ultra-cold freezer (-68 to -85°C).

---

**Shipping**

---



## TR-FRET TR beta Coactivator Assay Kit, goat

Dry Ice

---

### Size

800 x 20 µL assays

---

### Kit Components

The TR-FRET TR beta Coactivator Assay Kit contains TR beta-LBD (GST) protein, fluorescently labeled SRC2-2 coactivator peptide, Tb-labeled anti-GST antibody, and buffers. Store components as indicated in the assay protocol (-80°C, -20°C, or +4°C).

---

### Materials Required but Not Supplied

Microplate Reader

---

**Detection method** Fluorescent

---