



## BCL-2 TR-FRET Assay Kit

### Product Information

---

**Cat.No.**

Kit-1686

---

**Size**

384 reactions

---

**Description**

The BCL-2 TR-FRET Assay Kit is designed to measure the inhibition of BCL-2 binding to its ligand in a homogeneous 384 reaction format. This FRET-based assay requires no time-consuming washing steps, making it especially suitable for high throughput screening applications. The assay procedure is straightforward and simple; a sample containing terbiumlabeled donor, dye-labeled acceptor, BCL-2, peptide ligand, and an inhibitor is incubated for 120 minutes. Then, the fluorescence intensity is measured using a fluorescence reader.

---

**Applications**

Great for screening small molecular inhibitors for drug discovery and HTS applications.

---

**Storage**

At least 6 months from date of receipt when stored as directed. Kit components require different storage conditions. Be sure to store each component at the proper temperature upon arrival.

---

**Kit Components**

BCL-2: 10 µg; -80°C BCL-2 Peptide Ligand: 50 µl; -80°C Anti-His Tb donor: 2 x 10 µl; -20°C Dye-labeled acceptor: 2 x 10 µl; -20°C 3x BCL TR-FRET Assay Buffer: 4 ml; -20°C White, nonbinding, low volume, microtiter plate: 1; Room temp.

---