

# Active Recombinant Human ACE2 protein, His/Avi-tagged, Biotinylated

**Cat. No.** ACE2-051H    **Lot. No.** (See product label)

## SPECIFICATION

<b>Product Overview</b>	Biotinylated Recombinant Human ACE2(Gln18 - Ser740) protein, fused to His/Avi tag at the C-terminus, was expressed in CHO cells .
<b>Species</b>	Human
<b>Source</b>	CHO
<b>ProteinLength</b>	Gln18 - Ser740
<b>Description</b>	<p>Angiotensin I Converting Enzyme (ACE-2), also called ACEH (ACE homologue), is a dimeric, zinc-dependent metalloprotease of the ACE family that also includes somatic and germinal ACE (1, 2). ACE-2 mRNA is found at high levels in heart, testis, and kidney and at lower levels in a wide variety of tissues (1, 3). ACE-2 is the SARS-CoV and SARS-CoV2 Spike protein receptor in vivo (4-6), functions catalytically as a carboxypeptidase to cleave several substrates including angiotensins I and II, and acts as a partner for B0AT1-family amino acid transporters (1, 2). Through these functions, ACE-2 has been shown to be involved in several diseases including SARS, COVID19, acute lung injury (4, 7), heart disease (8), liver and lung fibrosis (9), inflammatory lung disease (10), and cardiopulmonary disease (11). Full length ACE-2 protein includes an extracellular region composed of a single N-terminal peptidase domain and C-terminal collectrin-like domain (CLD), a transmembrane domain, and a short cytoplasmic tail (12). The N-terminal peptidase region is required for binding to SARS-CoV and SARSCoV2 spike proteins, while the CLD contains a region that</p>

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

promotes dimerization and association with amino acid transporters (2). The peptidase domain contains a long deep cleft that undergoes a large hinge-bending movement at substrate and inhibitor binding (12). Classical ACE inhibitors such as captopril and lisinopril do not inhibit ACE-2 activity and inhibitors of ACE-2 do not inhibit ACE activity (13).


<b>Predicted N Terminal</b>	Protein identity confirmed by mass spectrometry.
<b>Form</b>	Supplied as a 0.2 µm filtered solution in Tris, NaCl, ZnCl <sub>2</sub> and Glycerol.
<b>Bio-activity</b>	Measured by its binding ability in a functional ELISA. When Recombinant SARS-CoV-2 Spike RBD Fc Chimera is immobilized at 0.2 g/mL (100 µ/well), Biotinylated Recombinant Human ACE2 His-tag Avi-tag binds with an ED50 of 1.2-9.6 ng/mL.
<b>Molecular Mass</b>	103-113 kDa, under reducing conditions
<b>Endotoxin</b>	<1.0 EU per 1 µg of the protein by the LAL method.
<b>Purity</b>	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
<b>Applications</b>	Bioactivity, Enzyme Activity
<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.6 months from date of receipt, -20 to -70 °C as supplied.3 months, -20 to -70 °C under sterile conditions after opening.
<b>Conjugation</b>	Biotin

## GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Gene Name</b>	ACE2 angiotensin I converting enzyme (peptidyl-dipeptidase A) 2 [ Homo sapiens ]
<b>Official Symbol</b>	ACE2
<b>Synonyms</b>	ACE2; angiotensin I converting enzyme (peptidyl-dipeptidase A) 2; angiotensin-converting enzyme 2; metalloprotease MPROT15; ACE-related carboxypeptidase; angiotensin I converting enzyme 2; angiotensin-converting enzyme homolog; ACEH;
<b>Gene ID</b>	59272
<b>mRNA Refseq</b>	NM_021804
<b>Protein Refseq</b>	NP_068576
<b>MIM</b>	300335
<b>UniProt ID</b>	Q9BYF1
<b>SDS-PAGE</b>	


Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127


45-1 Ramsey Road, Shirley, NY 11967, USA

2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.

### Binding Activity



### Enzyme Activity



Biotinylated Recombinant Human ACE2 His-tag Avi-tag(CHO Expressed) is measured by its ability to cleave a fluorogenic peptide substrate, Mca-YVADAPK(Dnp)-OH.

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA