

Recombinant Human ACVR1B cell lysate

Cat. No. ACVR1B-2130HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human ALK4 / ACVR1B derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[ACVR1B activin A receptor, type IB \[Homo sapiens \]](#)

Official Symbol

ACVR1B

Synonyms

ACVR1B; activin A receptor, type IB; ACVRLK4; activin receptor type-1B; ActRIB; ALK4; SKR2; activin receptor-like kinase 4; activin A receptor, type II-like kinase 4; serine/threonine-protein kinase receptor R2; ACTRIB;

Gene ID

[91](#)

mRNA Refseq

[NM_004302](#)

Protein Refseq


[NP_004293](#)

MIM


[601300](#)

UniProt ID

[P36896](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Chromosome
Location**

12q13

Pathway

Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Developmental Biology, organism-specific biosystem; MAPK signaling pathway, organism-specific biosystem; Regulation of Signaling by NODAL, organism-specific biosystem; Signaling by NODAL, organism-specific biosystem;

Function

ATP binding; SMAD binding; contributes_to activin binding; activin receptor activity, type I; activin receptor activity, type I; contributes_to activin-activated receptor activity; contributes_to growth factor binding; inhibin binding; metal ion binding; nucleotide binding; protein binding; protein serine/threonine kinase activity; protein serine/threonine kinase activity; receptor activity; transforming growth factor beta-activated receptor activity; transmembrane receptor protein serine/threonine kinase activity; ubiquitin protein ligase binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA