

Recombinant Human ADIPOQ cell lysate

Cat. No. ADIPOQ-1526HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human Adiponectin / Acrp30 / ADIPOQ derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

ADIPOQ adiponectin, C1Q and collagen domain containing [Homo sapiens]

Official Symbol

ADIPOQ

Synonyms

ADIPOQ; adiponectin, C1Q and collagen domain containing; ACDC, adipocyte, C1Q and collagen domain containing; adiponectin; ACRP30; AdipoQ; adipose most abundant gene transcript 1; apM1; GBP28; gelatin-binding protein 28; adipose specific collagen-like factor; 30 kDa adipocyte complement-related protein; adipocyte complement-related 30 kDa protein; adipose most abundant gene transcript 1 protein; ACDC; ADPN; APM1; APM-1; ADIPQTL1;

Gene ID

9370

mRNA Refseq

NM_001177800

Protein Refseq

NP_001171271

MIM

605441

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q15848
Chromosome Location	3q27
Pathway	Adipocytokine signaling pathway, organism-specific biosystem; Adipocytokine signaling pathway, conserved biosystem; Adipogenesis, organism-specific biosystem; Developmental Biology, organism-specific biosystem; PPAR signaling pathway, organism-specific biosystem; PPAR signaling pathway, conserved biosystem; Transcriptional Regulation of White Adipocyte Differentiation, organism-specific biosystem;
Function	cytokine activity; eukaryotic cell surface binding; hormone activity; identical protein binding; protein binding; protein homodimerization activity; receptor binding; sialic acid binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA