

# Recombinant Human Agouti Related Protein Homolog (Mouse), His-tagged

Cat. No. AGRP-136H Lot. No. (See product label)

## SPECIFICATION

<b>Product Overview</b>	Recombinant Human Agouti-related His-Tagged Fusion Protein, produced in <i>E. coli</i> , is 14.4 kDa (calculated) protein containing 112 amino acid residues of the human AGRP and 16 additional amino acid residues - His Tag, thrombin cleavage site. The AGRP is purified by proprietary chromatographic techniques.
<b>Species</b>	Human
<b>Source</b>	E.coli
<b>Description</b>	Agouti-related protein is an endogenous antagonist of hypothalamic alpha-melanocortin receptors MC3R and MC4R with potent orexigenic activity. Although a complete deletion of the AGRP gene does not produce any significant metabolic phenotypes, reduction in AGRP expression by RNA interference is associated with increased metabolic rate along with reduced weight gain. In hypothalamus, it is produced by neurons in the medial portion of arcuate nucleus, which produce also the potent orexigenic peptide Neuropeptide Y (NP-Y). Another site of central AGRP production is the hypothalamic nucleus. AGRP encompasses 132 amino acid residues and its alpha-melanocortin inhibiting activity results in a 34 amino acid cystine knot domain within the C-terminal (87-132) portion of the protein. Both AGRP and NP-Y expression was shown to be suppressed by leptin. Central administration of AGRP induces hyperphagia and increased gain in body weight in rodents, but may also exert metabolic effects even when hyperphagia is prevented. In the absence of hyperphagia, intracerebralventricular administration of AGRP caused significant

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

increases in plasma leptin and insulin concentrations (twofold and 1.5-fold, respectively) and fat pad mass. In the periphery, AGRP mRNA was found in adrenal glands, lung, testis, ovary, skeletal muscle and adipose tissue in humans or rodents. In the adrenals, it was shown that AGRP antagonizes glucocorticoid production mediated by MC4R. AGRP could then modulate locally the functions of some peripheral tissues such as adrenals. In human and rat serum, detectable levels of AGRP-like activity were reported in the lower picogram range. The serum AGRP levels were elevated in obese humans compared to lean controls and increased with fasting in rats.

**Amino Acid Sequence**

MKHHHHHHHM LVPRGSAQMG LAPMEGIRRP DQALLPELPG LGLRAPLKKT  
TAEQAEEDLL QEAQALAEVL DLQDREPRSS RRCVRLHESC LGQQVPCCDP  
CATCYCRFFN AFCYCRKLG T AMNPCSRT

**Physical Appearance**

Sterile Filtered White lyophilized (freeze-dried) powder.

**Form**

AGRP protein was lyophilized from 0.5 mg/ml in 5mM TRIS, 25mM NaCl, pH 7.5.

**Purity**

Purity of Agouti-related protein recombinant human is >95 % as determined by SDS-PAGE.

**Solubility**

Add deionized water to prepare a working stock solution of approximately 0.5 mg/ml and let the lyophilized pellet dissolve completely.

**Applications**

Western blotting, ELISA.

**Stability**

Store lyophilized AGRP protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C. The lyophilized protein remains stable until the expiry date when stored at -20°C.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

## GENE INFORMATION

<b>Gene Name</b>	AGRP agouti related protein homolog (mouse) [ Homo sapiens ]
<b>Synonyms</b>	AGRP; agouti related protein homolog (mouse); ART; AGRT; ASIP2; MGC118963; agouti related protein homolog; Agouti-related transcript, mouse, homolog of; Agouti-related protein
<b>Gene ID</b>	181
<b>mRNA Refseq</b>	NM_001138
<b>Protein Refseq</b>	NP_001129
<b>MIM</b>	602311
<b>UniProt ID</b>	O00253
<b>Chromosome Location</b>	16q22
<b>Pathway</b>	Adipocytokine signaling pathway; Neuroactive ligand-receptor interaction
<b>Function</b>	neuropeptide hormone activity; receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Agouti protein



Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA