

Recombinant Human AICDA 293 Cell Lysate

Cat. No. AICDA-8958HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for activation-induced cytidine deaminase (AICDA) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	AICDA activation-induced cytidine deaminase [Homo sapiens]
Official Symbol	AICDA
Synonyms	AICDA; activation-induced cytidine deaminase; AID; ARP2; CDA2; HIGM2; cytidine aminohydrolase; integrated into Burkitts lymphoma cell line Ramos;
Gene ID	57379
mRNA Refseq	NM_020661
Protein Refseq	NP_065712
MIM	605257
UniProt ID	Q9GZX7
Chromosome Location	12p13
Pathway	IL4-mediated signaling events, organism-specific biosystem; Intestinal immune network for IgA production, organism-specific biosystem; Intestinal immune network for IgA production, conserved biosystem; Primary immunodeficiency, organism-specific biosystem; Primary immunodeficiency, conserved biosystem; salvage pathways of pyrimidine deoxyribonucleotides, organism-specific biosystem; salvage pathways of pyrimidine ribonucleotides, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

cytidine deaminase activity; cytidine deaminase activity; hydrolase activity; hydrolase activity, acting on carbon-nitrogen (but not peptide) bonds, in cyclic amidines; metal ion binding; protein binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA