

## Recombinant Human ANAPC5 Protein, MYC/DDK-tagged

Cat. No. ANAPC5-2085H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant human ANAPC5 protein, fused to MYC/DDK-tagged at C-terminus, was expressed in HEK293
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	<p>This gene encodes a tetratricopeptide repeat-containing component of the anaphase promoting complex/cyclosome (APC/C), a large E3 ubiquitin ligase that controls cell cycle progression by targeting a number of cell cycle regulators such as B-type cyclins for 26S proteasome-mediated degradation through ubiquitination. The encoded protein is required for the proper ubiquitination function of APC/C and for the interaction of APC/C with transcription coactivators. It also interacts with polyA binding protein and represses internal ribosome entry site-mediated translation. Multiple transcript variants encoding different isoforms have been found for this gene. These differences cause translation initiation at a downstream AUG and result in a shorter protein (isoform b), compared to isoform a. [provided by RefSeq, Nov 2008].</p>
<b>Form</b>	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.
<b>Molecular Mass</b>	84.9 kDa
<b>Purity</b>	> 80% as determined by SDS-PAGE and Coomassie blue staining
<b>Concentration</b>	>50 ug/mL as determined by microplate BCA method

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



## GENE INFORMATION

**Gene Name** ANAPC5 anaphase promoting complex subunit 5 [ Homo sapiens ]

**Official Symbol** ANAPC5

**Synonyms** APC5; APC5

**Gene ID** 51433

**mRNA Refseq** NM\_001137559

**Protein Refseq** NP\_001131031

**MIM** 606948

**UniProt ID** Q9UJX4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA