

## Active Recombinant Human ANGPT2 protein, His-tagged

Cat. No. ANGPT2-531H Lot. No. (See product label)

### SPECIFICATION

**Product Overview** Active Recombinant Human ANGPT2 protein(O15123)(24~165 aa), fused with N-terminal His tag, was expressed in E.coli.

**Species** Human

**Source** E.coli

**ProteinLength** 24-165 aa

**Form** Lyophilized from sterile 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Bio-activity** Angiopoietin is part of a family of vascular growth factors that play a role in embryonic and postnatal angiogenesis. Angiopoietin cytokines are involved with controlling microvascular permeability, vasodilation, and vasoconstriction by signaling smooth muscle cells surrounding vessels. There are now four identified angiopoietins: ANGPT1, ANGPT2, ANGPT3, ANGPT4. Angiopoietin 2 (ANGPT2) promotes cell death and disrupts vascularization. Yet, when it is in conjunction with vascular endothelial growth factors, or VEGF, it can promote neo-vascularization. Besides, TEK Tyrosine Kinase (Tie2) has been identified as an interactor of ANGPT2, thus a binding ELISA assay was conducted to detect the interaction of recombinant human ANGPT2 and recombinant humanTie2. Briefly, ANGPT2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to Tie2-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

and incubated for 1h with anti-ANGPT2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µ stop solution to the wells and read at 450nm immediately. The binding activity of ANGPT2 and Tie2 was in a dose dependent manner.

<b>Molecular Mass</b>	17.6kDa
<b>Purity</b>	> 95% as determined by SDS-PAGE.
<b>Storage</b>	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
<b>Reconstitution</b>	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## GENE INFORMATION

<b>Gene Name</b>	ANGPT2 angiopoietin 2 [ Homo sapiens ]
<b>Official Symbol</b>	ANGPT2
<b>Synonyms</b>	ANGPT2; angiopoietin 2; angiopoietin-2; Ang2; ANG-2; Tie2-ligand; angiopoietin-2B; angiopoietin-2a; ANG2; AGPT2;
<b>Gene ID</b>	285
<b>mRNA Refseq</b>	NM_001118887
<b>Protein Refseq</b>	NP_001112359

 Tel: 1-631-559-9269 1-516-512-3133


 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


**MIM** 601922

**UniProt ID** O15123

**SDS-PAGE**



**Activity**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA