

Recombinant Human Angiopoietin-like 4, His-tagged

Cat. No. ANGPTL4-404H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human ANGPTL4 (aa 1-406) produced in <i>CHO cells</i> are fused at the C-terminus to a His-tag.
Species	Human
Source	Mammalian Cells
Protein Length	1-406 a.a.
Description	<p>The fasting-induced adipose factor (FIAF, ANGPTL4, PGAR, HFARP) was identified as an adipocytokine up-regulated by fasting, by peroxisome proliferator-activated receptor agonists, and by hypoxia. At the protein level, in human and mouse blood plasma, FIAF was found to be present both as a native protein and in a truncated form. Differentiation of mouse 3T3-L1 adipocytes was associated with the production of truncated FIAF, whereas in human white adipose tissue and SGBS adipocytes, only the native FIAF could be detected. Interestingly, the truncated FIAF was produced by human liver. Experimental data suggest that FIAF is mainly presented in human blood plasma in a truncated form (FIAF-S2), whose level is increased by fenofibrate treatment. Levels of both truncated and native FIAF showed marked inter individual variation but were not associated with body mass index and were not influenced by prolonged semistarvation.</p>
Source/Host	CHO cells.
Purity	≥90% (SDS-PAGE).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Formulation	Liquid. 0.2µm-filtered solution in PBS.
Endotoxin Content	<1EU/g protein (LAL-test).
Long Term Storage	-20°C.
Use/Stability	Working aliquots are stable for up to 3 months when stored at -20°C.
Handling	After opening, prepare aliquots and store at -20°C. Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	ANGPTL4 angiopoietin-like 4 [Homo sapiens]
Synonyms	ANGPTL4; angiopoietin-like 4; NL2; ARP4; FIAF; PGAR; HFARP; pp1158; ANGPTL2; fasting-induced adipose factor; PPARG angiopoietin related protein; hepatic angiopoietin-related protein; hepatic fibrinogen/angiopoietin-related protein; peroxisome proliferator-activated receptor (PPAR) gamma; induced angiopoietin-related protein; angiopoietin-related protein 4; PSEC0166, UNQ171/PRO197
Gene ID	51129
mRNA Refseq	NM_001039667
Protein Refseq	NP_001034756
MIM	605910
UniProt ID	Q9BY76
Chromosome	19p13.3

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Location

Pathway

PPAR signaling pathway; Metabolism of lipids and lipoproteins

Function

enzyme inhibitor activity; receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA