

## Recombinant Human ATF3 293 Cell Lysate

**Cat. No.** ATF3-8631HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for activating transcription factor 3 (ATF3), transcript variant 3 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

**Gene Name** ATF3 activating transcription factor 3 [ Homo sapiens ]

**Official Symbol** ATF3

**Synonyms** ATF3; activating transcription factor 3; cyclic AMP-dependent transcription factor ATF-3; cAMP-dependent transcription factor ATF-3; FLJ41705;

**Gene ID** 467

**mRNA Refseq** NM\_001030287

**Protein Refseq** NP\_001025458

**MIM** 603148

**UniProt ID** P18847

**Chromosome Location** 1q32.3

**Pathway** ATF-2 transcription factor network, organism-specific biosystem; Activation of Genes by ATF4, organism-specific biosystem; Diabetes pathways, organism-specific biosystem; Direct p53 effectors, organism-specific biosystem; Disease, organism-specific biosystem; HTLV-I infection, organism-specific biosystem; HTLV-I infection, conserved biosystem;

**Function** identical protein binding; protein binding; protein dimerization activity; sequence-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



specific DNA binding; sequence-specific DNA binding transcription factor activity;  
transcription corepressor activity;

☎ Tel: 1-631-559-9269 1-516-512-3133

☎ Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) ☎ Fax: 1-631-938-8127

☎ 45-1 Ramsey Road, Shirley, NY 11967, USA