

Recombinant Human ATP6V1C1 293 Cell Lysate

Cat. No. ATP6V1C1-8582HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for ATPase, H ⁺ transporting, lysosomal 42kDa, V1 subunit C1 (ATP6V1C1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	ATP6V1C1 ATPase, H+ transporting, lysosomal 42kDa, V1 subunit C1 [Homo sapiens]
Official Symbol	ATP6V1C1
Synonyms	ATP6C; ATP6D; ATP6V1C1; ATPase H+ transporting lysosomal (vacuolar proton pump) 42kD; ATPase H+ transporting lysosomal 42kD V1 subunit C isoform 1; ATPase H+ transporting lysosomal 42kDa V1 subunit C isoform 1; ATPase H+ transporting lysosomal 42kDa V1 subunit C1; ATPase H+ transporting lysosomal V1 subunit C1; FLJ20057; H(+) transporting two sector ATPase subunit C; H+ ATPase C subunit; H+ transporting ATPase chain C vacuolar; Subunit C of vacuolar proton ATPase V1 domain; V ATPase C subunit; V ATPase subunit C 1; V-ATPase subunit C 1; V-type proton ATPase subunit C 1; Vacuolar ATP synthase subunit C; Vacuolar proton pump 42 kD subunit; Vacuolar proton pump C subunit; Vacuolar proton pump subunit C 1; Vacuolar protonATPase subunit C VI domain; VATC; VATC1_HUMAN; VATPase C subunit; VATPase subunit C 1; VMA5; ATP6V1C1
Gene ID	528
mRNA Refseq	NM_001695.4
Protein Refseq	NP_001686.1
MIM	603097
UniProt ID	P21283

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	8q22.3
Pathway	Collecting duct acid secretion, organism-specific biosystem; Collecting duct acid secretion, conserved biosystem; Disease, organism-specific biosystem; Epithelial cell signaling in Helicobacter pylori infection, organism-specific biosystem; Epithelial cell signaling in Helicobacter pylori infection, conserved biosystem; Insulin receptor recycling, organism-specific biosystem; Iron uptake and transport, organism-specific biosystem
Function	hydrogen-exporting ATPase activity, phosphorylative mechanism; protein binding; proton-transporting ATPase activity, rotational mechanism; transporter activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA