

Active Recombinant Mouse Anxa5 protein, His-tagged

Cat. No. Anxa5-324M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Mouse Anxa5 protein(P48036)(1~319 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Mouse
Source	E.coli
ProteinLength	1-319 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Annexin V (ANXA5) is a multifunctional protein that is highly expressed on the apical surfaces of syncytiotrophoblasts, and plays an important role in haemostatic regulations, maintaining blood fluidity of the placenta. Lower ANXA5 levels have been observed in M2/ANXA5 haplotype carrying chorion. The association found between the maternal carriage of the M2/ANXA5 haplotype and an elevated risk of IUGR and/or PE supports the hypothesis that carrier status of this haplotype and the consequently reduced placental ANXA5 expression might be responsible, at least partially, for the onset of these gestational vascular complications. ANXA5 could be used as a biomarker for the early detection of PE and for the prediction of its severity. ANXA5 as an embryonic anticoagulant that appears deficient in contiguous specter of thrombophilia-related pregnancy complications culminating more frequently in miscarriage in a maternal M2 carrier background. Besides, Vinculin (VCL) has been

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

identified as an interactor of ANXA5, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse ANXA5 and recombinant human VCL. Briefly, ANXA5 were diluted serially in PBS, with 0.01% BSA (pH7.4). Duplicate samples of 100 µl were then transferred to VCL-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST and incubated for 1h with anti-ANXA5 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µ stop solution to the wells and read at 450 nm immediately. The binding activity of recombinant mouse ANXA5 and recombinant human VCL was shown, and this effect was in a dose dependent manner, the EC50 was 2.79 ug/ml.

Molecular Mass 37.0kDa

Endotoxin <1.0EU per 1µg (determined by the LAL method).

Purity > 95% as determined by SDS-PAGE.

Storage Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name [Anxa5 annexin A5 \[Mus musculus \]](#)

Official Symbol [Anxa5](#)

Synonyms ANXA5; annexin A5; PP4; CBP-I; PAP-I; VAC-alpha; annexin V; annexin-5; anchorin

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


CII; endonexin II; lipocortin V; calphobindin I; thromboplastin inhibitor; vascular anticoagulant-alpha; placental anticoagulant protein 4; placental anticoagulant protein I; Anx5; R74653;

Gene ID 11747


mRNA Refseq NM_009673

Protein Refseq NP_033803

SDS-PAGE



Activity



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA