

Recombinant Human ApoE2

Cat. No. ApoE2-3561H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	ApoE2
Species	Human
Source	E.coli
ProteinLength	317
Description	<p>Chylomicron remnants and very low density lipoprotein (VLDL) remnants are rapidly removed from the circulation by receptor-mediated endocytosis in the liver. Apolipoprotein E, a main apoprotein of the chylomicron, binds to a specific receptor on liver cells and peripheral cells. ApoE is essential for the normal catabolism of triglyceride-rich lipoprotein constituents. The APOE gene is mapped to chromosome 19 in a cluster with APOC1 and APOC2. Defects in apolipoprotein E result in familial dysbetalipoproteinemia, or type III hyperlipoproteinemia (HLP III), in which increased plasma cholesterol and triglycerides are the consequence of impaired clearance of chylomicron and VLDL remnants.</p>
Form	Lyophilized
Molecular Mass	36,154 Da
Purity	>90% by SDS-PAGE & HPLC analyses
Notes	Small volumes of ApoE2 recombinant protein may occasionally become entrapped in

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container's cap. Certain products may require to ship with dry ice.

Warning

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

GENE INFORMATION

Gene Name [ApoE2](#)

Official Symbol [ApoE2](#)

Synonyms [APOE](#); [APOE](#); [AD2](#); [LPG](#); [LDLCQ5](#); [Apo-E](#);

Gene ID [348](#)

mRNA Refseq [NM_000041.2](#)

Protein Refseq [NP_000032.1](#)

MIM [104310](#)

UniProt ID [P02649](#)

Chromosome Location [Chromosome: 19; NC_000019.10 \(44905782..44909393\)](#). Location: 19q13.2

Function

?ApoE is essential for the normal catabolism of triglyceride-rich lipoprotein constituents. The APOE gene is mapped to chromosome 19 in a cluster with APOC1 and APOC2. Defects in apolipoprotein E result in familial dysbetalipoproteinemia, or

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



type III hyperlipoproteinemia (HLP III), in which increased plasma cholesterol and triglycerides are the consequence of impaired clearance of chylomicron and VLDL remnants

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA