

Recombinant Human BACE1, C13&N15-labeled

Cat. No. BACE1-219H Lot. No. (See product label)

SPECIFICATION


Product Overview Recombinant Human BACE1 MS Standard Protein, C13 and N15-labeled (BACE1, Heavy Labeled) Thr 22 - Thr 457 (Accession # NP_036236.1) was produced in human 293 cells (HEK293).

Species Human

Source HEK293

ProteinLength 22-457 a.a.

Description Beta-secretase 1 (BACE1) also known as beta-site APP cleaving enzyme 1 (beta-site amyloid precursor protein cleaving enzyme 1), memapsin-2 (membrane-associated aspartic protease 2), and aspartyl protease 2 (ASP2), β -Secretase, and is a member of the peptidase A1 protein family, BACE1 is a type I integral membrane glycoprotein and aspartic protease that is found mainly in the Golgi. BACE1 is an aspartic-acid protease important in the pathogenesis of Alzheimer's disease, and in the formation of myelin sheaths in peripheral nerve cells.[1] The transmembrane protein contains two active site aspartate residues in its extracellular protein domain and may function as a dimer. This protease is responsible for the proteolytic processing of the amyloid precursor protein (APP). Generation of the 40 or 42 amino acid-long amyloid- β peptides that aggregate in the brain of Alzheimer's patients requires two sequential cleavages of the APP. Extracellular cleavage of APP by BACE creates a soluble extracellular fragment and a cell membrane-bound fragment referred to as C99. The elevation of BACE1 levels can be induced by amyloid plaques surrounding neurons at

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

early stages of pathology before neuron death occurs, and may drive a positive-feedback loop in AD. [2-3]

Form

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally Mannitol or Trehalose are added as protectants before lyophilization.

Molecular Mass

BACE1, Heavy Labeled, fused with 6×His tag at the C-terminus, has a calculated MW of 49.2 kDa. The predicted N-terminus is Thr 22. DTT-reduced Protein migrates as 50-65 kDa due to glycosylation.

Endotoxin

Less than 1.0 EU per µg of the BACE1, Heavy Labeled by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Storage

Avoid repeated freeze-thaw cycles.No activity loss was observed after storage at:In lyophilized state for 1 year (4oC-8oC); After reconstitution under sterile conditions for 1 month (4oC-8oC) or 3 months (-20oC to -70oC).

GENE INFORMATION

Gene Name

BACE1 beta-site APP-cleaving enzyme 1 [Homo sapiens]

Official Symbol

BACE1

Synonyms

BACE1; beta-site APP-cleaving enzyme 1; BACE, beta site APP cleaving enzyme; beta-secretase 1; asp 2; memapsin-2; APP beta-secretase; aspartyl protease 2; beta-site APP cleaving enzyme 1; beta-secretase 1 precursor variant 1; transmembrane aspartic proteinase Asp2; membrane-associated aspartic protease 2; beta-site amyloid beta A4 precursor protein-cleaving enzyme; ASP2; BACE; HSPC104; FLJ90568; KIAA1149;

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|----------------------------|---|
| Gene ID | 23621 |
| mRNA Refseq | NM_001207048 |
| Protein Refseq | NP_001193977 |
| MIM | 604252 |
| UniProt ID | P56817 |
| Chromosome Location | 11q23-q24 |
| Pathway | Alzheimers disease, organism-specific biosystem; Alzheimers disease, conserved biosystem; |
| Function | aspartic-type endopeptidase activity; beta-aspartyl-peptidase activity; enzyme binding; peptidase activity; |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA