

Recombinant Human basic leucine zipper transcription factor, ATF-like 3, His-tagged

Cat. No. BATF3-468H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	BATF3, 1-127aa Human, His tag, E.coli
Species	Human
Source	E.coli
ProteinLength	1-127 a.a.
Description	<p>Basic leucine zipper transcriptional factor ATF-like 3, also known as BATF3, is a 127 amino acid protein that localizes to the nucleus and contains one bZIP domain. Interacting with c-Jun, BATF3 functions as a negative regulator of AP-1-mediated transcription, specifically by heterodimerizing with c-Jun and binding to DNA response elements. Recombinant human BATF3 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.</p>
Form	Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.2M NaCl, 30% glycerol
Molecular Mass	16.9 kDa (150aa) confirmed by MALDI-TOF
AA Sequence	<p>MGSSHHHHHH SSGLVPRGSH MGSMSQGLPA AGSVLQRSVA APGNQPQPQP QQQSPEDDDR KVRREKNRV AAQRSRKKQT QKADKLHEEY ESLEQENTML RREIGKLTEE LKHLTEALKE HEKMCPLLLC PMNFVPVPPR PDPVAGCLPR</p>
Purity	>90% by SDS - PAGE

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Can be stored at +4C short term (1-2 weeks). For long term storage, aliquot and store at -20C or -70C. Avoid repeated freezing and thawing cycles.
Concentration	0.25 mg/ml (determined by BCA assay)
GENE INFORMATION	
Gene Name	BATF3 basic leucine zipper transcription factor, ATF-like 3 [Homo sapiens]
Official Symbol	BATF3
Synonyms	BATF3; basic leucine zipper transcription factor, ATF-like 3; basic leucine zipper transcriptional factor ATF-like 3; JDP1; Jun dimerization protein 1; JUNDM1; SNFT; B-ATF-3; Jun dimerization protein p21SNFT; 21-kD small nuclear factor isolated from T cells; 21 kDa small nuclear factor isolated from T-cells; FLJ36352; FLJ37535;
Gene ID	55509
mRNA Refseq	NM_018664
Protein Refseq	NP_061134
MIM	612470
UniProt ID	Q9NR55
Chromosome Location	1q32.3
Pathway	Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem; Calcium signaling in the CD4+ TCR pathway, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



biosystem;

Function

protein dimerization activity; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; transcription corepressor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA