

Recombinant Human BCAT1 293 Cell Lysate

Cat. No. BCAT1-8495HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for branched chain aminotransferase 1, cytosolic (BCAT1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	BCAT1 branched chain amino-acid transaminase 1, cytosolic [Homo sapiens]
Official Symbol	BCAT1
Synonyms	BCAT1; branched chain amino-acid transaminase 1, cytosolic; BCT1, branched chain aminotransferase 1, cytosolic; branched-chain-amino-acid aminotransferase, cytosolic; placental protein 18; branched chain aminotransferase 1, cytosolic; BCT1; PP18; BCATC; ECA39; MECA39; PNAS121; DKFZp686E12175;
Gene ID	586
mRNA Refseq	NM_001178091
Protein Refseq	NP_001171562
MIM	113520
UniProt ID	P54687
Chromosome Location	12pter-q12
Pathway	Branched-chain amino acid catabolism, organism-specific biosystem; Leucine degradation, leucine =>acetoacetate + acetyl-CoA, organism-specific biosystem; Leucine degradation, leucine => acetoacetate + acetyl-CoA, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

L-isoleucine transaminase activity; L-leucine transaminase activity; L-valine transaminase activity; branched-chain-amino-acid transaminase activity; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA