

Recombinant Human BIRC2 Protein (M1-S618), Tag Free

Cat. No. BIRC2-0713H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human GG-clAP1(M1-S618 end) (BIRC2) Protein was expressed in Insect cell.
Species	Human
Source	Insect Cells
ProteinLength	M1-S618
Description	<p>Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, mitogenic kinase signaling, and cell proliferation, as well as cell invasion and metastasis. Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and regulates both canonical and non-canonical NF-kappa-B signaling by acting in opposite directions: acts as a positive regulator of the canonical pathway and suppresses constitutive activation of non-canonical NF-kappa-B signaling. The target proteins for its E3 ubiquitin-protein ligase activity include: RIPK1, RIPK2, RIPK3, RIPK4, CASP3, CASP7, CASP8, TRAF2, DIABLO/SMAC, MAP3K14/NIK, MAP3K5/ASK1, IKBKG/NEMO, IKBKE and MXD1/MAD1. Can also function as an E3 ubiquitin-protein ligase of the NEDD8 conjugation pathway, targeting effector caspases for neddylation and inactivation. Acts as an important regulator of innate immune signaling via regulation of Toll-like receptors (TLRs), Nodlike receptors (NLRs) and RIG-I like receptors (RLRs), collectively referred to as pattern recognition receptors (PRRs). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

capability to kill cancer cells in a caspase-dependent and caspase-independent manner. Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8. Can stimulate the transcriptional activity of E2F1. Plays a role in the modulation of the cell cycle.

Form	Liquid
Endotoxin	< 0.01 EU per µg of the protein
Purity	90%
Stability	Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade.
Storage	Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	50 mM Tris-HCl (pH7.5), 200 mM NaCl, 20% glycerol
Shipping	It is shipped out with blue ice.

GENE INFORMATION

Gene Name	BIRC2 baculoviral IAP repeat containing 2 [Homo sapiens (human)]
Official Symbol	BIRC2
Synonyms	BIRC2; baculoviral IAP repeat containing 2; API1, baculoviral IAP repeat containing 2; baculoviral IAP repeat-containing protein 2; apoptosis inhibitor 1; c IAP1; cIAP1; hiap 2; MIHB; NFR2 TRAF signalling complex protein; RNF48; IAP-2; IAP homolog B; RING finger protein 48; inhibitor of apoptosis protein 2; baculoviral IAP repeat-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



containing 2; NFR2-TRAF signalling complex protein; TNFR2-TRAF-signaling complex protein 2; API1; HIAP2; Hiap-2; c-IAP1;

Gene ID 329

mRNA Refseq NM_001166

Protein Refseq NP_001157

MIM 601712

UniProt ID Q13490

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA