

GMP Recombinant Human BMP4

Cat. No. BMP4-125H Lot. No. (See product label)

SPECIFICATION

Product Overview	Disulfidelinked homodimer Mouse myeloma cell line, NS0 derived Ser293-Arg408 Accession # Q53XC5 Manufactured and tested under current Good Manufacturing Practice (GMP) guidelines.
Species	Human
Source	NS0
Description	BMPs are secreted signaling molecules that comprise a subfamily of the TGF-beta superfamily and were originally identified as regulators of cartilage and bone formation. There are at least 20 structurally and functionally related BMPs, most of which play roles in embryogenesis and morphogenesis of various tissues and organs. Biologically active BMPs are usually homodimers containing a characteristic cysteine knot structure. Heterodimers, BMP-2/BMP-7 and BMP-4/BMP-7 have also been suggested to exist and function in vivo. They are more potent inducers of bone formation than their respective homodimers. In addition, heterodimers, but not homodimers, are ventral mesoderm inducers. Heterodimer activity may be mediated by a different or additional receptor subtype.
Predicted N Terminal	Ser293-Pro-Lys-His-His-Ser-Gln-Arg-Ala-Arg
Form	Lyophilized from a 0.2 µm filtered solution in Acetonitrile and TFA.
Bio-activity	Measured by its ability to induce alkaline phosphatase production by ATDC5 mouse chondrogenic cells. Binnerts, M.E. et al. (2004) Biochem. Biophys. Res. Commun.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

315(2):272. The ED50 for this effect is typically 2.5-15 ng/mL.

Molecular Mass

Predicted Molecular Mass: 13 kDa (monomer).SDSPAGE: 22-25 kDa, reducing conditions; 37-41 kDa, nonreducing conditions

Endotoxin

<0.01 EU per 1 µg of the protein by the LAL method.

Purity

>95%, by SDSPAGE with silver staining, under reducing conditions.

Storage

Use a manual defrost freezer and avoid repeated freezethaw cycles. 12 months, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.

Reconstitution

Reconstitute at 50-200 µg/mL in 4 mM HCl.

GENE INFORMATION

Gene Name

[BMP4 bone morphogenetic protein 4 \[Homo sapiens \]](#)

Official Symbol

[BMP4](#)

Synonyms

BMP4; bone morphogenetic protein 4; BMP2B; BMP-4; BMP-2B; bone morphogenetic protein 2B; ZYME; OFC11; BMP2B1; MCOPS6;

Gene ID

[652](#)

mRNA Refseq

[NM_001202](#)

Protein Refseq

[NP_001193](#)

MIM

[112262](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q53XC5
Chromosome Location	14q22-q23
Pathway	Adipogenesis, organism-specific biosystem; Basal cell carcinoma, organism-specific biosystem; Basal cell carcinoma, conserved biosystem; Glypican 3 network, organism-specific biosystem; Heart Development, organism-specific biosystem; Hedgehog signaling pathway, organism-specific biosystem; Hedgehog signaling pathway, conserved biosystem;
Function	BMP receptor binding; chemoattractant activity; cytokine activity; growth factor activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA