

Recombinant Human BTG2 293 Cell Lysate

Cat. No. BTG2-8392HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for BTG family, member 2 (BTG2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	BTG2 BTG family, member 2 [Homo sapiens]
Official Symbol	BTG2
Synonyms	BTG2; BTG family, member 2; protein BTG2; B cell translocation gene 2; MGC126063; MGC126064; nerve growth factor inducible anti proliferative; NGF inducible anti proliferative protein PC3; PC3; pheochromacytoma cell 3; TIS21; pheochromacytoma cell-3; B-cell translocation gene 2; NGF-inducible anti-proliferative protein PC3; nerve growth factor-inducible anti-proliferative;
Gene ID	7832
mRNA Refseq	NM_006763
Protein Refseq	NP_006754
MIM	601597
UniProt ID	P78543
Chromosome Location	1q32
Pathway	Direct p53 effectors, organism-specific biosystem; RNA degradation, organism-specific biosystem; RNA degradation, conserved biosystem;
Function	RNA polymerase II core promoter proximal region sequence-specific DNA binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



transcription factor activity involved in positive regulation of transcription;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA