

Recombinant Human BTK cell lysate

Cat. No. BTK-001HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human Bruton Tyrosine Kinase / BTK Kinase derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[BTK Bruton agammaglobulinemia tyrosine kinase \[Homo sapiens \]](#)

Official Symbol

BTK

Synonyms

BTK; Bruton agammaglobulinemia tyrosine kinase; AGMX1, IMD1; tyrosine-protein kinase BTK; ATK; PSCTK1; XLA; B-cell progenitor kinase; agammaglobulinaemia tyrosine kinase; tyrosine-protein kinase BTK isoform (lacking exon 14); dominant-negative kinase-deficient Brutons tyrosine kinase; AT; BPK; IMD1; AGMX1; MGC126261; MGC126262;

Gene ID

[695](#)

mRNA Refseq

[NM_000061](#)

Protein Refseq

[NP_000052](#)

MIM

[300300](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q06187
Chromosome Location	Xq21.33-q22
Pathway	Adaptive Immune System, organism-specific biosystem; Antigen Activates B Cell Receptor Leading to Generation of Second Messengers, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; B cell receptor signaling pathway, organism-specific biosystem; B cell receptor signaling pathway, conserved biosystem; BCR signaling pathway, organism-specific biosystem; Class I PI3K signaling events, organism-specific biosystem;
Function	ATP binding; ATP binding; identical protein binding; kinase activity; lipid binding; metal ion binding; non-membrane spanning protein tyrosine kinase activity; nucleotide binding; phosphatidylinositol-3,4,5-trisphosphate binding; protein binding; protein kinase activity; protein tyrosine kinase activity; protein tyrosine kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA