

# Recombinant Human BTLA Protein, hFc-tagged, FITC conjugated

Cat. No. BTLA-233HF Lot. No. (See product label)

## SPECIFICATION

<b>Product Overview</b>	FITC conjugated recombinant human BTLA (Lys31-Ser150) protein was fused to human IgG1 Fc tag at C-terminus and expressed in human 293 cells (HEK293).
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	Lys31-Ser150
<b>Description</b>	B- and T-lymphocyte attenuator (BTLA) is also known as B- and T-lymphocyte-associated protein, CD antigen CD272. BTLA contains one Ig-like V-type (immunoglobulin-like) domain. As a lymphocyte inhibitory receptor, BTLA/CD272 inhibits lymphocytes during immune response. BTLA/CD272 can interact with tyrosine phosphatases PTPN6/SHP-1 and PTPN11/SHP-2, and interact with TNFRSF14/HVEM.
<b>Form</b>	Lyophilized
<b>Molecular Mass</b>	The protein has a calculated MW of 40.4 kDa. The protein migrates as 50-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
<b>N-terminal Sequence Analysis</b>	Lys 31

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Endotoxin</b>	< 1.0 EU/ µg by the LAL method.
<b>Purity</b>	> 92 % as determined by SDS-PAGE
<b>Characteristic</b>	<p>Disulfide-linked homodimer</p> <p>Labeled with FITC via amines</p> <p>Excitation source: 488 nm spectral line, argon-ion laser</p> <p>Excitation Wavelength: 488 nm</p> <p>Emission Wavelength: 535 nm</p>
<b>Storage</b>	<p>For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower.</p> <p>Please avoid repeated freeze-thaw cycles.</p> <p>This product is stable after storage at:</p> <p>-20 to -70 centigrade for 12 months in lyophilized state;</p> <p>-70 centigrade for 3 months under sterile conditions after reconstitution.</p>
<b>Storage Buffer</b>	Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.
<b>Conjugation</b>	FITC

## GENE INFORMATION

<b>Gene Name</b>	<b>BTLA</b>
<b>Official Symbol</b>	<b>BTLA</b>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



<b>Synonyms</b>	BTLA; B and T lymphocyte associated; B- and T-lymphocyte attenuator; BTLA1; CD272; B and T lymphocyte attenuator; B- and T-lymphocyte-associated protein; FLJ16065; MGC129743
<b>Gene ID</b>	151888
<b>mRNA Refseq</b>	NM_001085357
<b>Protein Refseq</b>	NP_001078826
<b>MIM</b>	607925
<b>UniProt ID</b>	Q7Z6A9

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA