

Recombinant Human CAMK2B cell lysate

Cat. No. CAMK2B-596HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human CAMK2B / CaMKII-beta derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

CAMK2B calcium/calmodulin-dependent protein kinase II beta [Homo sapiens]

Official Symbol

CAMK2B

Synonyms

CAMK2B; calcium/calmodulin-dependent protein kinase II beta; calcium/calmodulin dependent protein kinase (CaM kinase) II beta , CAMKB; calcium/calmodulin-dependent protein kinase type II subunit beta; calcium/calmodulin dependent protein kinase type II beta chain; CaM kinase II beta subunit; CaM kinase II beta chain; CAM2; CAMK2; proline rich calmodulin dependent protein kinase; caMK-II subunit beta; CaM-kinase II beta chain; proline rich calmodulin-dependent protein kinase; CAMKB; MGC29528;

Gene ID

816

mRNA Refseq

NM_001220

Protein Refseq

NP_001211

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	607707
UniProt ID	Q13554
Chromosome Location	7p14.3-p14.1
Pathway	Activation of NMDA receptor upon glutamate binding and postsynaptic events, organism-specific biosystem; Amphetamine addiction, organism-specific biosystem; Amphetamine addiction, conserved biosystem; CREB phosphorylation through the activation of CaMKII, organism-specific biosystem; CREB phosphorylation through the activation of Ras, organism-specific biosystem; Calcium Regulation in the Cardiac Cell, organism-specific biosystem; Calcium signaling pathway, organism-specific biosystem;
Function	ATP binding; actin binding; calmodulin binding; calmodulin-dependent protein kinase activity; nucleotide binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA