

Recombinant Human CASR 293 Cell Lysate

Cat. No. CASR-7825HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for calcium-sensing receptor (CASR) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CASR calcium-sensing receptor [Homo sapiens]
Official Symbol	CASR
Synonyms	CASR; calcium-sensing receptor; HHC, HHC1, hypocalciuric hypercalcemia 1; extracellular calcium-sensing receptor; FHH; GPRC2A; NSHPT; severe neonatal hyperparathyroidism; parathyroid Ca(2+)-sensing receptor 1; parathyroid cell calcium-sensing receptor; CAR; FIH; HHC; EIG8; HHC1; PCAR1; MGC138441;
Gene ID	846
mRNA Refseq	NM_000388
Protein Refseq	NP_000379
UniProt ID	P41180
Chromosome Location	3q21.1
Pathway	Class C/3 (Metabotropic glutamate/pheromone receptors), organism-specific biosystem; E-cadherin signaling in keratinocytes, organism-specific biosystem; G alpha (i) signalling events, organism-specific biosystem; G alpha (q) signalling events, organism-specific biosystem; GPCR downstream signaling, organism-specific biosystem; GPCR ligand binding, organism-specific biosystem; GPCRs, Class C Metabotropic glutamate, pheromone, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

G-protein coupled receptor activity; phosphatidylinositol phospholipase C activity; protein binding; receptor activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA