

Recombinant Human CBR3 293 Cell Lysate

Cat. No. CBR3-7810HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for carbonyl reductase 3 (CBR3) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CBR3 carbonyl reductase 3 [Homo sapiens]
Official Symbol	CBR3
Synonyms	CBR3; carbonyl reductase 3; carbonyl reductase [NADPH] 3; SDR21C2; short chain dehydrogenase/reductase family 21C; member 2; carbonyl reductase (NADPH) 3; NADPH-dependent carbonyl reductase 3; short chain dehydrogenase/reductase family 21C, member 2; hCBR3;
Gene ID	874
mRNA Refseq	NM_001236
Protein Refseq	NP_001227
MIM	603608
UniProt ID	O75828
Chromosome Location	21q22.2
Pathway	Arachidonic acid metabolism, organism-specific biosystem; Arachidonic acid metabolism, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism of xenobiotics by cytochrome P450, organism-specific biosystem; Metabolism of xenobiotics by cytochrome P450, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

3-keto sterol reductase activity; NADPH binding; carbonyl reductase (NADPH) activity; nucleotide binding; oxidoreductase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA