

## Active Recombinant Human Chemokine (C-C Motif) Ligand 13, MlgG2a Fc-tagged

Cat. No. CCL13-276H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	The extracellular domain of human CCL13 (AAH08621.1)(Gln24-Thr98) is fused to the N-terminus of the Fc region of mouse IgG2a was expressed in CHO cell.
<b>Species</b>	Human
<b>Source</b>	CHO
<b>ProteinLength</b>	24-98 a.a.
<b>Description</b>	This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into the arterial wall during atherosclerosis.
<b>Form</b>	Lyophilized from 0.2µm-filtered solution in PBS.
<b>Bio-activity</b>	Measured by its ability to chemoattract 2-day cultured human monocytes.
<b>Molecular Mass</b>	35KDa (monomer)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>AA Sequence</b>	Gln24-Thr98
<b>Endotoxin</b>	<0.06 eu/μg="" as="" determined="" by="" lal="">
<b>Purity</b>	>98%, by SDS-PAGE under reducing conditions.
<b>Stability</b>	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
<b>Reconstitution</b>	Reconstitute at 100μg/ml in sterile PBS.
<b>Warning</b>	Avoid freeze/thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">CCL13 chemokine (C-C motif) ligand 13 [ Homo sapiens ]</a>
<b>Official Symbol</b>	CCL13
<b>Synonyms</b>	CCL13; chemokine (C-C motif) ligand 13; SCYA13, small inducible cytokine subfamily A (Cys Cys), member 13; C-C motif chemokine 13; CKb10; MCP 4; MGC17134; NCC 1; SCYL1; CK-beta-10; new CC chemokine 1; small-inducible cytokine A13; monocyte chemotactic protein 4; monocyte chemoattractant protein 4; small inducible cytokine subfamily A (Cys-Cys), member 13; NCC1; MCP-4; NCC-1; SCYA13;
<b>Gene ID</b>	<a href="#">6357</a>
<b>mRNA Refseq</b>	<a href="#">NM_005408</a>
<b>Protein Refseq</b>	<a href="#">NP_005399</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



<b>MIM</b>	601391
<b>UniProt ID</b>	Q99616
<b>Chromosome Location</b>	17q11.2
<b>Pathway</b>	Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;
<b>Function</b>	chemokine activity; receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA