

Active Recombinant Human Chemokine (C-C Motif) Ligand 13, HIgG1 Fc-tagged

Cat. No. CCL13-278H Lot. No. (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CCL13 (AAH08621.1)(Gln24-Thr98) is fused to the N-terminus of the Fc region of human IgG1 was expressed in CHO cell
Species	Human
Source	CHO
ProteinLength	24-98 a.a.
Description	<p>This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into the arterial wall during atherosclerosis.</p>
Form	Lyophilized from 0.2µm-filtered solution in PBS.
Bio-activity	Measured by its ability to chemoattract 2-day cultured human monocytes.
Molecular Mass	35KDa (monomer)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	Gln24-Thr98
Endotoxin	<0.06 eu/μg="" as="" determined="" by="" lal="" test="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100μg/ml in sterile PBS.
Warning	Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	CCL13 chemokine (C-C motif) ligand 13 [Homo sapiens]
Official Symbol	CCL13
Synonyms	CCL13; chemokine (C-C motif) ligand 13; SCYA13, small inducible cytokine subfamily A (Cys Cys), member 13; C-C motif chemokine 13; CKb10; MCP 4; MGC17134; NCC 1; SCYL1; CK-beta-10; new CC chemokine 1; small-inducible cytokine A13; monocyte chemotactic protein 4; monocyte chemoattractant protein 4; small inducible cytokine subfamily A (Cys-Cys), member 13; NCC1; MCP-4; NCC-1; SCYA13;
Gene ID	6357
mRNA Refseq	NM_005408
Protein Refseq	NP_005399

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



MIM	601391
UniProt ID	Q99616
Chromosome Location	17q11.2
Pathway	Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;
Function	chemokine activity; receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA