

Recombinant Human Chemokine (C-C Motif) Ligand 13, His-tagged

Cat. No. CCL13-2792H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human CCL13 protein (24-98aa), fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Species	Human
Source	E.coli
ProteinLength	24-98 a.a.
Description	CCL13 is a small cytokine belonging to the CC chemokine family that is also known as thymus and activation regulated chemokine (TARC). This protein induces chemotaxis in monocytes, eosinophils, T lymphocytes, and basophils by binding cell surface G-protein linked chemokine receptors such as CCR2, CCR3 and CCR5. It can be induced by the inflammatory cytokines interleukin-1 and TNF- α .
Sequences	MGSSHHHHHH SSGLVPRGSH MQPDALNVPS TCCFTFSSKK ISLQRLKSYV ITTSRCPQKA VIFRTKLGKE ICADPKEKWV QNYMKHLGRK AHTLKT.
Form	Liquid. In 20 mM Tris-HCl buffer (pH7.4) containing 0.1M NaCl, 1mM DTT, 10% glycerol.
Molecular Weight	10.8 kDa (96aa), confirmed by MALDI-TOF.
Purity	> 90% by SDS – PAGE.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Concentration 0.25 mg/ml (determined by Bradford assay).

Storage Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

GENE INFORMATION

Gene Name [CCL13 chemokine \(C-C motif\) ligand 13 \[Homo sapiens \]](#)

Synonyms

CCL13; chemokine (C-C motif) ligand 13; C-C motif chemokine 13; CK-beta-10; monocyte chemoattractant protein 4; monocyte chemotactic protein 4; new CC chemokine 1; small inducible cytokine subfamily A (Cys-Cys), member 13; small-inducible cytokine A13; NCC1; CKb10; MCP-4; NCC-1; SCYL1; SCYA13; MGC17134

Gene ID [6357](#)

mRNA Refseq [NM_005408](#)

Protein Refseq [NP_005399](#)

MIM [601391](#)

UniProt ID [Q99616](#)

Chromosome Location 17q11.2

Pathway Chemokine signaling pathway; Cytokine-cytokine receptor interaction; NOD-like receptor signaling pathway

Function chemokine activity; receptor binding; signal transducer activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA