

Active Recombinant Human CCL15 Protein (92 aa)

Cat. No. CCL15-070C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CCL15 Protein without tag was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	92
Description	<p>CCL15, a new human CC chemokine, was isolated from a human fetal spleen cDNA library. CCL15 cDNA encodes a predicted 113 amino acid (aa) protein containing a putative signal peptide of 21 amino acids that is cleaved to generate a 92 aa residue mature protein. Within the CC family members, human CCL15 shares 45%, 44%, 35%, and 30% aa homology with mouse C10, human MIP-1, human HCC-1, and mouse MIP-1γ, respectively. The gene for MIP-5 is found on chromosome 17 where the genes for most of the human CC chemokines are located. Human CCL15 is expressed in T and B lymphocytes, NK cells, monocytes and monocyte-derived dendritic cells. Human MIP-5 is chemotactic for T cells and monocytes and has been shown to induce calcium flux in human CCR-1-transfected cells.</p>
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	Fully biologically active when compared to standard. Determined by its ability to chemoattract human T lymphocytes using a concentration range of 1.0 -10.0 ng/mL, corresponding to a Specific Activity of 1×10^5 IU/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	10.1 kDa, a single non-glycosylated polypeptide chain containing 92 amino acids.
AA Sequence	QFTNDAETELMMSKLPLENPVVLNSFHFAADCCTSYISQSIPCSLMKSYFETSSECS KPGVIFLTKKGRQVCAKPSGPGVQDCMKKLPYSI
Endotoxin	Less than 1 EU/mg of rHuMIP-5/CCL15 as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analyses.
Storage	This lyophilized preparation is stable at 2-8 centigrade, but should be kept at -20 centigrade for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 centigrade. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 to -70 centigrade. Avoid repeated freeze/thaw cycles.
Storage Buffer	Lyophilized from a 0.2mm filtered concentrated solution in 20mM PB, pH 7.4, 100mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	CCL15 chemokine (C-C motif) ligand 15 [Homo sapiens]
Official Symbol	CCL15
Synonyms	CCL15; chemokine (C-C motif) ligand 15; SCYA15, small inducible cytokine subfamily

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

A (Cys Cys), member 15; C-C motif chemokine 15; CC chemokine 3; chemokine CC 2; HCC 2; HMRP 2B; leukotactin 1; Lkn 1; macrophage inflammatory protein 5; MIP 1 delta; MIP 1d; MIP 5; NCC 3; SCYL3; MIP-1 delta; chemokine CC-2; new CC chemokine 3; small-inducible cytokine A15; small inducible cytokine subfamily A (Cys-Cys), member 15; LKN1; NCC3; SY15; HCC-2; LKN-1; MIP-5; NCC-3; MIP-1D; MRP-2B; SCYA15; HMRP-2B;

Gene ID [6359](#)

mRNA Refseq [NM_032965](#)

Protein Refseq [NP_116741](#)

MIM [601393](#)

UniProt ID [Q16663](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA