

Recombinant Human CCL17 protein

Cat. No. CCL17-24H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CCL17 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	71
Description	Human CCL17 also known as thymus and activation-related chemokine (TARC) is encoded by the CCL17 gene located on the chromosome 16 in humans. It is expressed by thymus cells constitutively and phytohemagglutinin-stimulated peripheral blood mononuclear cells transiently. CCL17 signals through the chemokine receptors CCR4 and CCR8 and displays chemotactic activity for T lymphocytes and some other leukocytes. It plays an important role in skin diseases such as atopic dermatitis, bullous pemphigoid and mycosis fungoides. CCL17 has approximately 24 – 29 % amino acid sequence identity with RANTES, MIP-1 α , MIP-1 β , MCP-1, MCP-2, MCP-3 and I-309.
Form	Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM PB, pH 7.4, 150mM NaCl.
Bio-activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 1.0-10 ng/ml.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 8.1 kDa, a single non-glycosylated polypeptide chain containing 71 amino acids.
AA Sequence	ARGTNVGRECCLEYFKGAIPLRKLKTWYQTSEDCSRDAIVFVTVQGRAICSDPNNKR VKNAVKYLQSLERS
Endotoxin	Less than 1 EU/μg of rHuTARC/CCL17 as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	CCL17
Official Symbol	CCL17
Synonyms	CCL17; chemokine (C-C motif) ligand 17; SCYA17, small inducible cytokine subfamily A (Cys Cys), member 17; C-C motif chemokine 17; ABCD 2; TARC; CC chemokine TARC; T cell-directed CC chemokine; small-inducible cytokine A17; thymus and activation-regulated chemokine; small inducible cytokine subfamily A (Cys-Cys),

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



member 17; ABCD-2; SCYA17; A-152E5.3; MGC138271; MGC138273;

Gene ID [6361](#)

mRNA Refseq [NM_002987](#)

Protein Refseq [NP_002978](#)

MIM [601520](#)

UniProt ID [Q92583](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA