

Active Recombinant Human Chemokine (C-C Motif) Ligand 18 (Pulmonary And Activation-Regulated)

Cat. No. CCL18-300H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human Macrophage Inflammatory Protein-4 produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 69 amino acids and having a molecular mass of 7813 Daltons.
Species	Human
Source	E.coli
Description	Chemokine (C-C motif) ligand 18 (CCL18) is a small cytokine belonging to the CC chemokine family that was previously called PARC (pulmonary and activation-regulated chemokine).
Form	Lyophilized with no additives.
Purity	Purity Greater than 98% as determined by: Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.
Bio-activity	Human Macrophage Inflammatory Protein-4 is fully biologically active when compared to standard. The Activity is calculated by the ability to chemoattract Human T lymphocytes at 1.0-8.0 ng/ml.
Physical Appearance	Sterile Filtered white lyophilized (freeze-dried) powder.
Solubility	Centrifuge vial before opening. When reconstituting the product, gently pipet and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/ml, which can be further diluted into other aqueous solutions.

Purity Greater than 98% as determined by

Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.

Protein Content determined by

UV spectroscopy at 280 nm. Analysis by RP-HPLC calibrated against a known standard. Quantitation on SDS-PAGE against a known standard.

Endotoxin Level

Endotoxin level as measured by LAL is <0.01ng/ug or <0.1EU/ug.

Amino acid sequence

AQVGTNKELC CLVYTSWQIP QKFLVDYSET SPQCPKPGVI LLTKRGRQIC
ADPNKKWVQK YISDLKLNA

Storage

Lyophilized product is very stable at -20 degrees. Reconstituted material should be aliquoted and frozen at -20. It is recommended to add a carrier protein (0.1% HSA or BSA) for long term storage.

GENE INFORMATION

Gene Name

CCL18 chemokine (C-C motif) ligand 18 (pulmonary and activation-regulated) [Homo sapiens]

Official Symbol

CCL18

Synonyms

CKb7; PARC; AMAC1; DCCK1; MIP-4; AMAC-1; DC-CK1; SCYA18;
CCL18_HUMAN; C-C motif chemokine 18 [Precursor]; Small-inducible cytokine A18; Macrophage inflammatory protein 4; pulmonary and activation-regulated

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

chemokine;CC chemokine PARC;Alternative macrophage activation-associated CC chemokine 1; Dendritic cell chemokine 1; CCL18; chemokine (C-C motif) ligand 18 (pulmonary and activation-regulated)

Gene ID [6362](#)

mRNA Refseq [NM_002988](#)

Protein Refseq [NP_002979](#)

MIM [603757](#)

UniProt ID [P55774](#)

Chromosome Location 17q11.2

Pathway chemokine activity

Function Cytokine-cytokine receptor interaction

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA