

# Active Recombinant Human Chemokine (C-C motif) Ligand 19, HIgG1 Fc-tagged

Cat. No. CCL19-163H Lot. No. (See product label)

## SPECIFICATION

<b>Product Overview</b>	The extracellular domain of human CCL19 (AAH27968.1)(Gly22-Ser98) is fused to the N-terminus of the Fc region of human IgG1 was expressed in CHO cell
<b>Species</b>	Human
<b>Source</b>	CHO
<b>ProteinLength</b>	22-98 a.a.
<b>Description</b>	This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene may play a role in normal lymphocyte recirculation and homing. It also plays an important role in trafficking of T cells in thymus, and in T cell and B cell migration to secondary lymphoid organs. It specifically binds to chemokine receptor CCR7.
<b>Form</b>	Lyophilized from 0.2µm-filtered solution in PBS.
<b>Bio-activity</b>	Measured by its ability to chemoattract 5-10 day cultured human peripheral blood lymphocytes (PBL).
<b>Molecular Mass</b>	35KDa (monomer)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>AA Sequence</b>	Gly22-Ser98
<b>Endotoxin</b>	<0.06 eu/μg="" as="" determined="" by="" lal="">
<b>Purity</b>	>98%, by SDS-PAGE under reducing conditions.
<b>Stability</b>	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
<b>Reconstitution</b>	Reconstitute at 100μg/ml in sterile PBS.
<b>Warning</b>	Avoid freeze/thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">CCL19 chemokine (C-C motif) ligand 19 [ Homo sapiens ]</a>
<b>Official Symbol</b>	CCL19
<b>Synonyms</b>	CCL19; chemokine (C-C motif) ligand 19; SCYA19, small inducible cytokine subfamily A (Cys Cys), member 19; C-C motif chemokine 19; beta chemokine exodus 3; CC chemokine ligand 19; CK beta 11; CKb11; EBI1 ligand chemokine; ELC; exodus 3; macrophage inflammatory protein 3 beta; MIP 3b; exodus-3; CK beta-11; MIP-3-beta; EBI1-ligand chemokine; beta chemokine exodus-3; beta-chemokine exodus-3; small-inducible cytokine A19; macrophage inflammatory protein 3-beta; epstein-Barr virus-induced molecule 1 ligand chemokine; small inducible cytokine subfamily A (Cys-Cys), member 19; MIP3B; MIP-3b; SCYA19; MGC34433;
<b>Gene ID</b>	<a href="#">6363</a>
<b>mRNA Refseq</b>	<a href="#">NM_006274</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Protein Refseq</b>	NP_006265
<b>MIM</b>	602227
<b>UniProt ID</b>	Q99731
<b>Chromosome Location</b>	9p13
<b>Pathway</b>	Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; G alpha (i) signalling events, organism-specific biosystem;
<b>Function</b>	CCR chemokine receptor binding; CCR10 chemokine receptor binding; CCR7 chemokine receptor binding; chemokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA