

Recombinant Human CCL2 Protein, Biotinylated

Cat. No. CCL2-016H Lot. No. (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant human CCL2 protein was tag free expressed.
Species	Human
ProteinLength	99
Description	<p>This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Chemokines are a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. Elevated expression of the encoded protein is associated with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection.</p>
Form	Lyophilized
AA Sequence	<p>MKVSAALLCLLLIAATFIPQGLAQPDAINAPVTCCYNFTNRKISVQRLASYRRITSSKC PKEAVIFKTIVAKEICADPKQKWVQDSMDHLDKQTQTPKT</p>
Purity	> 97%

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered and lyophilized.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.
Conjugation	Biotin

GENE INFORMATION

Gene Name	CCL2 chemokine (C-C motif) ligand 2 [Homo sapiens (human)]
Official Symbol	CCL2
Synonyms	CCL2; chemokine (C-C motif) ligand 2; SCYA2, small inducible cytokine A2 (monocyte chemotactic protein 1, homologous to mouse Sig je); C-C motif chemokine 2; GDCF 2; HC11; MCAF; MCP 1; MCP1; MGC9434; monocyte chemoattractant protein 1; monocyte chemotactic and activating factor; monocyte chemotactic protein 1; homologous to mouse Sig je; monocyte secretory protein JE; small inducible cytokine subfamily A (Cys Cys); member 2; SMC CF; small-inducible cytokine A2; monocyte chemoattractant protein-1; small inducible cytokine subfamily A (Cys-Cys), member 2; small inducible cytokine A2 (monocyte chemotactic protein 1, homologous

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



to mouse Sig-je); MCP-1; SCYA2; GDCF-2; SMC-CF; HSMCR30;

Gene ID [6347](#)

mRNA Refseq [NM_002982](#)

Protein Refseq [NP_002973](#)

MIM [158105](#)

UniProt ID [P13500](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA