

Recombinant Human CCL4 protein

Cat. No. CCL4-294H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CCL4 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	69
Description	CCL4 encoded by the CCL4 gene, also known as Macrophage Inflammatory Protein-1 β (MIP-1 β) is a CC chemokine with specificity for CCR5 receptors and it is a major HIV-suppressive factor produced by CD8+ T cells. It is a monokine with inflammatory and chemokinetic properties. Recombinant CCL4 induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). Specifically, MIP-1-beta (3-69) is also a ligand for CCR1 and CCR2 isoform B. Recombinant human CCL4 contains 69 amino acids and it shares 77 % and 80 % a.a. sequence identity with murine and rat CCL4, respectively. Both human and murine MIP-1 α and MIP-1 β are active on human and murine hematopoietic cells.
Form	Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Bio-activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood monocytes is in a concentration range of 5.0-20 ng/ml.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 7.8 kDa, a single non-glycosylated polypeptide chain containing 69 amino acids.
AA Sequence	APMGSDPPTACCFSTARKLPRNFVVDYYETSSLCSQPAVVFQTKRSKQVCADPSE SWVQEYVYDLELN
Endotoxin	Less than 1 EU/μg of rHuMIP-1β/CCL4 as determined by LAL method.
Purity	>96% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	CCL4
Official Symbol	CCL4
Synonyms	CCL4; chemokine (C-C motif) ligand 4; LAG1, SCYA4, small inducible cytokine A4 (homologous to mouse Mip 1b); C-C motif chemokine 4; Act 2; AT744.1; MIP 1 beta; PAT 744; SIS-gamma; MIP-1-beta(1-69); CC chemokine ligand 4; secreted protein G-26; T-cell activation protein 2; small-inducible cytokine A4; lymphocyte-activation

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



gene 1; G-26 T-lymphocyte-secreted protein; lymphocyte activation gene 1 protein; macrophage inflammatory protein 1-beta; small inducible cytokine A4 (homologous to mouse Mip-1b); ACT2; G-26; HC21; LAG1; LAG-1; MIP1B; SCYA2; SCYA4; MIP1B1; MIP-1-beta; MGC104418; MGC126025; MGC126026;

Gene ID 6351

mRNA Refseq NM_002984

Protein Refseq NP_002975

MIM 182284

UniProt ID P13236

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA