

## Recombinant Human CCL4L1 293 Cell Lysate

**Cat. No.** CCL4L1-7720HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for chemokine (C-C motif) ligand 4-like 1 (CCL4L1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

**Gene Name** CCL4L1 chemokine (C-C motif) ligand 4-like 1 [ Homo sapiens ]

**Official Symbol** CCL4L1

**Synonyms** CCL4L1; chemokine (C-C motif) ligand 4-like 1; CCL4L, chemokine (C C motif) ligand 4 like , SCYA4L, small inducible cytokine A4 like; C-C motif chemokine 4-like; AT744.2; LAG 1; MIP-1-beta; macrophage inflammatory protein-1b2; lymphocyte activation gene 1 protein; macrophage inflammatory protein 1-beta; monocyte adherence-induced protein 5-alpha; LAG1; CCL4L; LAG-1; SCYA4L;

**Gene ID** 9560

**mRNA Refseq** NM\_001001435

**Protein Refseq** NP\_001001435

**MIM** 603782

**UniProt ID** Q8NHW4

**Chromosome Location** 17q12

**Pathway** Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Cytosolic DNA-sensing pathway, organism-specific biosystem; Cytosolic DNA-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



sensing pathway, conserved biosystem; IL12-mediated signaling events, organism-specific biosystem;

**Function**

chemokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA