

## Active Human CCL5

**Cat. No.** CCL5-114H    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	It is a synthetic Protein
<b>Species</b>	Human
<b>Description</b>	<p>RANTES (acronym for Regulated upon Activation, Normal T cell Expressed and presumably Secreted), was initially discovered by subtractive hybridization as a transcript expressed in T cells but not B cells. Eosinophil-chemotactic activities released by thrombin-stimulated human platelets have also been purified and found to be identical to RANTES. Besides T cells and platelets, RANTES has been reported to be produced by renal tubular epithelium, synovial fibroblasts and selected tumor cells. Human RANTES cDNA encodes a highly basic 91 amino acid residue precursor polypeptide with a 23 amino acid residue hydrophobic signal peptide that is cleaved to generate the 68 amino acid residue mature protein. Murine RANTES has also been cloned. Human and mouse RANTES share approximately 85% sequence homology at the amino acid level and exhibit cross-species activities.</p>
<b>Form</b>	Lyophilized
<b>Bio-activity</b>	Activity of this protein measured by its monocyte chemotactic activity and by its ability to chemoattract human CCR5 transfected mouse BaF/3 cells. ED50's for these activities are typically 0.01-0.03 µg/ml and 1-5 ng/ml respectively.
<b>Applications</b>	Activity Assay
<b>Storage</b>	Product should be stored at -20°C. Aliquot to avoid freeze/thaw cycles

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Reconstitution** Recommended reconstitution in sterile phosphate buffered saline which contains at least 0.1% human or bovine serum albumin to prepare a stocksolution of no less than 20 µg/ml and vortex thoroughly.

## GENE INFORMATION

**Gene Name** CCL5 chemokine (C-C motif) ligand 5 [ Homo sapiens ]

**Official Symbol** CCL5

**Synonyms**

CCL5; chemokine (C-C motif) ligand 5; D17S136E, SCYA5, small inducible cytokine A5 (RANTES); C-C motif chemokine 5; beta chemokine RANTES; MGC17164; RANTES; regulated upon activation; normally T expressed; and presumably secreted; SIS delta; SISd; small inducible cytokine subfamily A (Cys Cys); member 5; T cell specific protein p288; T cell specific RANTES protein; TCP228; eoCP; SIS-delta; beta-chemokine RANTES; small-inducible cytokine A5; T-cell specific protein p288; t cell-specific protein P228; T-cell-specific protein RANTES; eosinophil chemotactic cytokine; small inducible cytokine A5 (RANTES); small inducible cytokine subfamily A (Cys-Cys), member 5; regulated upon activation, normally T-expressed, and presumably secreted; SCYA5; D17S136E;

**Gene ID** 6352

**mRNA Refseq** NM\_002985

**Protein Refseq** NP\_002976

**MIM** 187011

**UniProt ID** P13501

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Chromosome Location</b>	17q11.2-q12
<b>Pathway</b>	Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem;
<b>Function</b>	CCR1 chemokine receptor binding; CCR1 chemokine receptor binding; CCR1 chemokine receptor binding; CCR4 chemokine receptor binding; CCR5 chemokine receptor binding; chemoattractant activity; chemokine activity; chemokine activity; chemokine receptor antag

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA