

Recombinant Human CCL8

Cat. No. CCL8-28723TH **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant full length MCP2 protein , is a non-glycosylated, polypeptide chain containing 76 amino acids and has a MW of 8.9 kDa.
Species	Human
Source	E.coli
Description	This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils. By recruiting leukocytes to sites of inflammation this cytokine may contribute to tumor-associated leukocyte infiltration and to the antiviral state against HIV infection.
Biological activity	The biological activity was determined by measuring the dose dependent mobilization of intracellular calcium (calcium flux) with human THP-1 cells. Significant calcium mobilization is observed with >500ng/mL of recombinant human MCP2. Human MCP2 also indu
Form	Lyophilised:Lyophilized from a concentrated (1mg/ml) sterile solution containing no additives. Reconstitute the lyophilized product in sterile 18MO-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	>95% by SDS-PAGE
Storage buffer	Sterile solution containing no additives.
Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
Sequences of amino acids	The sequence of the first five N-terminal amino acids was determined and was found to be Gln-Pro-Asp-Ser-Val.
Full Length	Full L.

GENE INFORMATION

Gene Name	CCL8 chemokine (C-C motif) ligand 8 [Homo sapiens]
Official Symbol	CCL8
Synonyms	CCL8; chemokine (C-C motif) ligand 8; SCYA8, small inducible cytokine subfamily A (Cys Cys), member 8 (monocyte chemotactic protein 2); C-C motif chemokine 8; HC14; MCP 2;
Gene ID	6355
mRNA Refseq	NM_005623
Protein Refseq	NP_005614
MIM	602283
Uniprot ID	P80075


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Chromosome Location	17q11.2
Pathway	Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;
Function	chemokine activity; heparin binding; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA