

Recombinant Human CCR1 293 Cell Lysate

Cat. No. CCR1-7698HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for chemokine (C-C motif) receptor 1 (CCR1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CCR1 chemokine (C-C motif) receptor 1 [Homo sapiens]
Official Symbol	CCR1
Synonyms	CCR1; chemokine (C-C motif) receptor 1; CMKBR1, SCYAR1; C-C chemokine receptor type 1; CD191; CKR 1; MIP1aR; CCR-1; CC-CKR-1; RANTES-R; C-C CKR-1; MIP-1alpha-R; LD78 receptor; RANTES receptor; macrophage inflammatory protein 1-alpha receptor; CKR1; CKR-1; HM145; CMKBR1; SCYAR1;
Gene ID	1230
mRNA Refseq	NM_001295
Protein Refseq	NP_001286
MIM	601159
UniProt ID	P32246
Chromosome Location	3p21
Pathway	Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; G alpha (i) signalling

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



events, organism-specific biosystem;

Function

C-C chemokine binding; C-C chemokine receptor activity; G-protein coupled receptor activity; chemokine (C-C motif) ligand 5 binding; chemokine (C-C motif) ligand 7 binding; chemokine receptor activity; phosphatidylinositol phospholipase C activity; protein binding; receptor activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA