

Human CCR5 peptide

Cat. No. CCR5-366H Lot. No. (See product label)

SPECIFICATION

Product Overview CCR5 peptide is used for blocking the activity of Anti-CCR5

Species Human

Description

This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. This protein is expressed by T cells and macrophages, and is known to be an important co-receptor for macrophage-tropic virus, including HIV, to enter host cells. Defective alleles of this gene have been associated with the HIV infection resistance. The ligands of this receptor include monocyte chemoattractant protein 2 (MCP-2), macrophage inflammatory protein 1 alpha (MIP-1 alpha), macrophage inflammatory protein 1 beta (MIP-1 beta) and regulated on activation normal T expressed and secreted protein (RANTES). Expression of this gene was also detected in a promyeloblastic cell line, suggesting that this protein may play a role in granulocyte lineage proliferation and differentiation. This gene is located at the chemokine receptor gene cluster region. Two transcript variants encoding the same protein have been found for this gene.

Form

Liquid in PBS, pH 7.2 (10 mM NaH₂PO₄ + 130 mM NaCl) containing 0.1% bovine serum albumin and 0.02% sodium azide. Precaution: Sodium azide is a poisonous and hazardous substance which should be handled by trained staff only.

Applications

Incubating the peptide with equal volume of antibody for 30 min at 37°C usually completely blocks the activity of Anti-CCR5 in Western blot.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	When stored at -20°C, stable for one year from date of receipt.
Concentration	0.2 mg/ml
GENE INFORMATION	
Gene Name	CCR5 chemokine (C-C motif) receptor 5 (gene/pseudogene) [Homo sapiens(human)]
Official Symbol	CCR5
Synonyms	CCR5; CKR5; CCR-5; CD195; CKR-5; CCCKR5; CMKBR5; IDDM22; CC-CKR-5; chemokine (C-C motif) receptor 5 (gene/pseudogene); C-C chemokine receptor type 5; chemr13; HIV-1 fusion coreceptor; chemokine receptor CCR5; C-C motif chemokine receptor 5 A159A
Gene ID	1234
mRNA Refseq	NM_000579
Protein Refseq	NP_000570
MIM	601373
UniProt ID	P51681
Chromosome Location	3p21.31
Pathway	Binding and entry of HIV virion; Chemokine receptors bind chemokines; Chemokine signaling pathway

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

C-C chemokine binding; actin binding; chemokine (C-C motif) ligand 5 binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA