

Recombinant Human CCT2 293 Cell Lysate

Cat. No. CCT2-7691HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for chaperonin containing TCP1, subunit 2 (beta) (CCT2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CCT2 chaperonin containing TCP1, subunit 2 (beta) [Homo sapiens]
Official Symbol	CCT2
Synonyms	CCT2; chaperonin containing TCP1, subunit 2 (beta); T-complex protein 1 subunit beta; Cctb; T-complex protein 1, beta subunit; chaperonin containing t-complex polypeptide 1, subunit 2; chaperonin containing t-complex polypeptide 1, beta subunit; CCTB; 99D8.1; PRO1633; CCT-beta; TCP-1-beta; MGC142074; MGC142076;
Gene ID	10576
mRNA Refseq	NM_001198842
Protein Refseq	NP_001185771
MIM	605139
UniProt ID	P78371
Chromosome Location	12q14
Pathway	Association of TriC/CCT with target proteins during biosynthesis, organism-specific biosystem; Chaperonin-mediated protein folding, organism-specific biosystem; Cooperation of Prefoldin and TriC/CCTin actin and tubulin folding, organism-specific biosystem; Folding of actin by CCT/TriC, organism-specific biosystem; Formation of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



tubulin folding intermediates by CCT/TriC, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; Prefoldin mediated transfer of substrate to CCT/TriC, organism-specific biosystem;

Function

ATP binding; nucleotide binding; unfolded protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA