

Active Recombinant Human CD19 Protein, Fc-tagged, Atto 488 conjugated

Cat. No. CD19-07H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human Atto 488 conjugated CD19 (Glu21-Lys291) protein with a human IgG1 (Pro100-Lys330) Fc tag at C-terminus was expressed in chinese hamster ovary cell line.
Species	Human
Source	CHO
ProteinLength	21-291 a.a.
Description	<p>This gene encodes a member of the immunoglobulin gene superfamily. Expression of this cell surface protein is restricted to B cell lymphocytes. This protein is a reliable marker for pre-B cells but its expression diminishes during terminal B cell differentiation in antibody secreting plasma cells. The protein has two N-terminal extracellular Ig-like domains separated by a non-Ig-like domain, a hydrophobic transmembrane domain, and a large C-terminal cytoplasmic domain. This protein forms a complex with several membrane proteins including complement receptor type 2 (CD21) and tetraspanin (CD81) and this complex reduces the threshold for antigen-initiated B cell activation. Activation of this B-cell antigen receptor complex activates the phosphatidylinositol 3-kinase signalling pathway and the subsequent release of intracellular stores of calcium ions. This protein is a target of chimeric antigen receptor (CAR) T-cells used in the treatment of lymphoblastic leukemia. Mutations in this gene are associated with the disease common variable immunodeficiency 3</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

(CVID3) which results in a failure of B-cell differentiation and impaired secretion of immunoglobulins. CVID3 is characterized by hypogammaglobulinemia, an inability to mount an antibody response to antigen, and recurrent bacterial infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Form	<p>Disulfide-linked homodimer</p> <p>Labeled with Atto 488 via amines</p> <p>Excitation Wavelength: 501 nm</p> <p>Emission Wavelength: 523 nm</p>
Bio-activity	<p>Measured by flow cytometry for its ability to bind anti-Human CD19 Monoclonal Antibody conjugated fluorescent beads.</p>
Molecular Mass	57 kDa
N-terminal Sequence Analysis	Glu21
Endotoxin	<1.0 EU/μg of the protein by the LAL method.
Purity	>90%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Storage	<p>Protect from light. Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <p>6 months from date of receipt, -20 to -70 centigrade as supplied.</p> <p>1 month, 2 to 8 centigrade under sterile conditions after opening.</p> <p>3 months, -20 to -70 centigrade under sterile conditions after opening.</p>
Storage Buffer	Supplied as a 0.2 μm filtered solution in PBS with BSA as a carrier protein.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Shipping The product is shipped with dry ice or equivalent.

Conjugation Atto 488

GENE INFORMATION

Gene Name CD19 CD19 molecule [Homo sapiens (human)]

Official Symbol CD19

Synonyms CD19; CD19 molecule; B4; CVID3; B-lymphocyte antigen CD19; B-lymphocyte surface antigen B4; T-cell surface antigen Leu-12; differentiation antigen CD19

Gene ID 930

mRNA Refseq NM_001770

Protein Refseq NP_001761

MIM 107265


UniProt ID P15391

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

FC




Human PBMC CD4+ and CD8+ T cells were either (A) transduced or (B) not transduced (negative control) with a Human CD19 CAR and then cultured for 11 days. Cells were washed twice with 2 mL 1x PBS then stained with a viability dye for 15-30 min at room temp in the dark. Human block was added during the viability stain to block Fc receptors. Cells were washed twice with Flow Cytometry Staining Buffer then stained with Human CD4 PE-Cy7-conjugated Antibody and rhCD19 Atto 488-conjugated protein for 25 min at room temp in the dark. Staining was performed using our staining membrane proteins protocol.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE

2 µg/lane of Recombinant Human CD19 Fc Chimera Atto 488 was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 77-92 kD and 154-184 kDa, respectively.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA