

CD19 protein-coupled magnetic MicroBeads

Cat. No. CD19-5078M **Lot. No.** (See product label)

SPECIFICATION


Species Mouse

Capacity for 2×10^9 total cells

Background The mouse monoclonal antibody reacts with mouse CD19, which is expressed on B cell lineage cells, throughout their development from early pro-B cell stages on, and is down-regulated upon terminal differentiation to plasma cells. Mouse CD19 MicroBeads are used for positive selection or depletion of mouse B cells from various tissues such spleen.

Application Positive selection or depletion of CD19+ B cells spleen.
Isolation of CD19+ B cells to study the accessibility of enhancer elements during B cell development by in vivo footprinting.
Isolation of pure B cells for preparation of cell extracts to analyze conditions for V(D)J recombination in a cell-free system.

Analysis



A single-cell suspension from mouse spleen was prepared using the gentleMACS Dissociator. CD19+ B cells were isolated from this single-cell suspension using the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



CD19 MicroBeads, and autoMACS Pro Separator. Cells were fluorescently stained with REAfinity Antibodies CD45R (B220)-APC-Vio 770 (clone REA755), mouse, and CD19-PE-Vio770, mouse (clone REA749) and analyzed by flow cytometry using the MACSQuant Analyzer. Viable leukocytes were gated for analysis based on scatter signals and 7-AAD Staining Solution fluorescence.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA