

Recombinant Human CD1D cell lysate

Cat. No. CD1D-2479HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human CD1D / R3G1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

CD1D CD1d molecule [[Homo sapiens](#)]

Official Symbol

CD1D

Synonyms

CD1D; CD1d molecule; CD1d antigen , CD1D antigen, d polypeptide; antigen-presenting glycoprotein CD1d; R3G1; thymocyte antigen CD1D; CD1D antigen, d polypeptide; T-cell surface glycoprotein CD1d; differentiation antigen CD1-alpha-3; HMC class I antigen-like glycoprotein CD1D; R3; CD1A; MGC34622;

Gene ID

912

mRNA Refseq

[NM_001766](#)

Protein Refseq

[NP_001757](#)

MIM

188410

UniProt ID

[P15813](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Chromosome Location	1q22-q23
Pathway	Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem; Hematopoietic cell lineage, organism-specific biosystem; Hematopoietic cell lineage, conserved biosystem;
Function	beta-2-microglobulin binding; cell adhesion molecule binding; eukaryotic cell surface binding; exogenous lipid antigen binding; histone binding; lipid antigen binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA