

Recombinant Human CD247 Protein, C-His-tagged

Cat. No. CD247-171H Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant Human CD247 Protein with C-His tag was expressed in E. coli.

Species Human

Source E.coli

Description

When T cells encounter antigens via the T cell receptor (TCR), information about the quantity and quality of antigens is relayed to the intracellular signal transduction machinery. This activation process depends mainly on CD3 (Cluster of Differentiation 3), a multiunit protein complex that directly associates with the TCR. CD3 is composed of four polypeptides: ζ , γ , ϵ and δ . Each of these polypeptides contains at least one immunoreceptor tyrosine-based activation motif (ITAM). Engagement of TCR complex with foreign antigens induces tyrosine phosphorylation in the ITAM motifs and phosphorylated ITAMs function as docking sites for signaling molecules such as ZAP-70 and p85 subunit of PI-3 kinase. TCR ligation also induces a conformational change in CD3 ϵ , such that a proline region is exposed and then associates with the adaptor protein Nck. The CD3 ζ invariant chain is a type-I transmembrane protein that exists in the TCR signaling complex as a disulfide-linked homodimer. The cytoplasmic tail of each CD3 ζ monomer contains three distinct ITAM motifs, each containing two tyrosine residues. Phosphorylation of CD3 ζ ITAM tyrosine residues, including Y142, is driven by recruitment of the Lck and Fyn tyrosine kinases to the TCR. Lck/Fyn-mediated ITAM phosphorylation creates docking sites that promote the SH2 domain-dependent recruitment and activation of ZAP-70, which drives amplification of signaling events downstream of the TCR that facilitate T cell

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

activation. Phosphorylation of a pool of p16 CD3 ζ leads to the generation of p21 and p23 species, which differ in the degree of ITAM phosphorylation. It has been proposed that the ratio of p21/p23 contributes to regulating the amplitude of T cell activation. CD3 ζ plays an important role in the assembly and surface expression of the TCR complex. Indeed, research studies have demonstrated that CD3 ζ is degraded in response to Ag-dependent TCR stimulation as a mechanism to tightly control T cell activation.

Molecular Mass ~12 kDa

AA Sequence RVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPQRRKN
PQEGLYNELQKDKMAEAYSEIGMKGERRRGKGGHDGLYQGLSTATKDTYDALHMQA
LPPR

Purity Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Notes For research use only, not for use in diagnostic procedure.

Storage Store at 4 centigrade short term. Aliquot and store at -20 centigrade long term. Avoid freeze-thaw cycles.

Concentration ≥ 0.5 mg/mL

Storage Buffer PBS, 4M Urea, pH7.4

GENE INFORMATION

Gene Name CD247 CD247 molecule [Homo sapiens (human)]

Official Symbol CD247

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms CD247; CD247 molecule; CD3Z, CD3z antigen, zeta polypeptide (TiT3 complex) , CD247 antigen; T-cell surface glycoprotein CD3 zeta chain; CD3H; CD3Q; CD3zeta chain; TCR zeta chain; CD247 antigen, zeta subunit; T-cell receptor T3 zeta chain; CD3Z antigen, zeta polypeptide (TiT3 complex); T-cell antigen receptor complex, zeta subunit of CD3; T3Z; CD3Z; TCRZ; CD3-ZETA;

Gene ID 919


mRNA Refseq NM_000734

Protein Refseq NP_000725

MIM 186780

UniProt ID P20963

SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA