

Active Recombinant Human CD27, MlgG2a Fc-tagged

Cat. No. CD27-001H Lot. No. (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CD27[TNFRSF7] (AAH12160.1)(Thr21- Ile192) is fused to the N-terminus of the Fc region of mouse IgG2a was expressed in CHO cell
Species	Human
Source	CHO
Description	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.
Form	Lyophilized from 0.2µm-filtered solution in PBS.
Bio-activity	Measured by its ability to inhibit the proliferation of mouse T cells induced by CD27 Ligand and anti-CD3.
Molecular Mass	45 KDa (monomer)
AA Sequence	Thr21- Ile192

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	<0.06 eu/μg="" as="" determined="" by="" lal="" test.="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100μg/ml in sterile PBS.
Warning	Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	CD27 CD27 molecule [Homo sapiens]
Official Symbol	CD27
Synonyms	CD27; CD27 molecule; TNFRSF7, tumor necrosis factor receptor superfamily, member 7; CD27 antigen; S152; Tp55; CD27L receptor; T cell activation antigen S152; T-cell activation antigen CD27; tumor necrosis factor receptor superfamily, member 7; T14; TNFRSF7; MGC20393;
Gene ID	939
mRNA Refseq	NM_001242
Protein Refseq	NP_001233
MIM	186711
UniProt ID	P26842

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Chromosome Location	12p13
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;
Function	cysteine-type endopeptidase inhibitor activity involved in apoptotic process; protein binding; receptor activity; transmembrane signaling receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA