

Recombinant Human CD27 molecule Protein, Tag Free

Cat. No. CD27-229H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human CD27/TNFRSF7 protein (P26842) (Ala 20-Arg 191) without tag was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	20-191aa
Description	<p>The tumor necrosis factor (TNF) receptor family is composed of several type I integral membrane glycoproteins that exhibit homology in their cysteine-rich extracellular domains. Members of this family include FAS, OX40, CD27 and CD30. Ligands for these receptors are often type II transmembrane glycoproteins, as is the case for CD27 and CD30. CD27 is a homodimeric lymphocyte-specific surface antigen present on T and B lymphocytes. Activation of the CD3 complex via the T cell receptor for antigen leads to an increase in CD27 expression. Together, CD27 and its ligand, CD27L, generate co-stimulatory signals required for complete T cell activation. CD30 is a surface marker for neoplastic cells of the Hodgkin's lymphoma and related hematologic malignancies. CD30L has been shown to enhance the proliferation of the Hodgkin's cell line HDLM-2, but exerts antiproliferative effects on large cell anaplastic lymphoma cell lines.</p>
Tag	Non
Molecular Mass	19.6 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 1.0 EU/μg by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade. It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage Buffer	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.
Reconstitution	Reconstitute at 250 μg/mL in sterile water.

GENE INFORMATION

Gene Name	CD27 CD27 molecule [Homo sapiens (human)]
Official Symbol	CD27
Synonyms	CD27; CD27 molecule; TNFRSF7, tumor necrosis factor receptor superfamily, member 7; CD27 antigen; S152; Tp55; CD27L receptor; T cell activation antigen S152; T-cell activation antigen CD27; tumor necrosis factor receptor superfamily, member 7; T14; TNFRSF7; MGC20393;
Gene ID	939
mRNA Refseq	NM_001242
Protein Refseq	NP_001233
MIM	186711

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID

P26842

Protein on SDS-
PAGE under
reducing (R)
condition.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA